

# Can one use total variation prior for edge-preserving Bayesian inversion?

Matti Lassas<sup>1</sup> and Samuli Siltanen<sup>2</sup>

<sup>1</sup> Institute of Mathematics, Helsinki University of Technology, PO Box 1100, FIN-02015 TKK, Finland

<sup>2</sup> GE Healthcare, PO Box 20, FIN-04301 Tuusula, Finland

Received 16 October 2003, in final form 19 May 2004

Published 6 August 2004

Online at [stacks.iop.org/IP/20/1537](http://stacks.iop.org/IP/20/1537)

doi:10.1088/0266-5611/20/5/013

## Abstract

Estimation of non-discrete physical quantities from indirect linear measurements is considered. Bayesian solution of such an inverse problem involves discretizing the problem and expressing available *a priori* information in the form of a prior distribution in a finite-dimensional space. Since *a priori* information is independent of the measurement, the discretization of the unknown quantity can be arbitrarily fine regardless of the number of measurements. The main result is that Bayesian conditional mean estimates for total variation prior distribution are not edge-preserving with very fine discretizations of the model space. Theoretical findings are illustrated by a numerical example with computer simulated data.

## 1. Introduction

Consider an indirect noisy measurement  $m$  of a physical quantity  $u$ :

$$m = Au + \varepsilon, \quad (1.1)$$

where  $\varepsilon$  is random noise and the linear operator  $A$  models the measurement. The corresponding inverse problem is

$$\text{given } m, \quad \text{find } u. \quad (1.2)$$

We assume that the object  $u$  is *a priori* known to be piecewise regular. Our aim is to raise methodological concerns about the solution of (1.2) with Bayesian inversion using total variation prior distribution.

Practical solution of (1.2) with Bayesian inversion requires discretization of the problem and expressing available *a priori* information on  $u$  in the form of a *prior distribution*  $\pi_{\text{pr}}^{(n)}$  in an  $n$ -dimensional subspace  $Y_n \subset Y$ , where  $u$  is *a priori* known to belong to some function space  $Y$ . Let  $m$  and  $\varepsilon$  be random vectors taking values in  $\mathbb{R}^N$ , and denote their distributions

by  $\pi_m$  and  $\pi_\varepsilon$ , respectively. Given a realization  $\hat{m}$  of the measurement  $m = Au + \varepsilon$ , Bayes' formula yields the *posterior distribution* for the random variable  $u_n$  taking values in  $Y_n$ :

$$\pi(u_n | \hat{m}) = \frac{\pi_{\text{pr}}^{(n)}(u_n)\pi(\hat{m} | u_n)}{\pi_m(\hat{m})} \sim \pi_{\text{pr}}^{(n)}(u_n)\pi_\varepsilon(\hat{m} - Au_n). \quad (1.3)$$

The approximate solution of (1.2) is given as some point estimate for (1.3). Such estimates include the *maximum a posteriori* (MAP) and *conditional mean* (CM) estimates defined by

$$u_n^{\text{MAP}} = \arg \max_{u_n \in Y_n} \pi(u_n | \hat{m}), \quad u_n^{\text{CM}} = \int_{Y_n} u_n \pi(u_n | \hat{m}) \, du_n,$$

respectively. The crucial step in Bayesian inversion is the construction of  $Y_n$  and  $\pi_{\text{pr}}^{(n)}$ . The probability measure  $\pi_{\text{pr}}^{(n)}$  should assign high probability to functions  $u_n \in Y_n$  that are typical in light of *a priori* information on  $u$ , and low probability to atypical functions. For more details, see [12, 15, 17, 22, 25, 31].

*Total variation (TV) regularization* for edge-preserving noise removal was introduced in [29]. This deterministic inversion method was later successfully applied to other inverse problems where ‘blocky’ reconstructions are desired [9, 35]. TV regularization is equivalent to determining the MAP estimate for (1.3) with TV prior distribution. This observation inspired consideration of CM and other Bayesian estimates using the TV prior [17, 19, 30]; preservation of edges was achieved with fixed discretization.

However, from the pure Bayesian point of view, *a priori* information and its discrete representation are independent of the measurement, and the dimension  $n$  can be freely chosen. In our view, the space  $Y_n$  and the distribution  $\pi_{\text{pr}}^{(n)}$  should be constructed for all  $n$  so that the following two conditions are satisfied:

- (i) There is a random function  $v$  taking values in  $Y$  such that  $\lim_{n \rightarrow \infty} u_n = v$ . This guarantees that the representation of *a priori* information becomes more accurate when  $n$  grows.
- (ii) There are continuous linear operators  $T_n : Y \rightarrow Y_n$  such that  $u_n = T_n v$ . This means that the finite-dimensional approximations to  $u$  are achieved systematically from the limit function  $v$ . (We restrict ourselves to linear discretizations  $T_n$  as we consider linear inverse problems only.)

Any choice of  $Y_n$  and  $\pi_{\text{pr}}^{(n)}$  having properties (i) and (ii) is called *discretization invariant*.

We show (for a generic one-dimensional problem) that the TV prior is not discretization invariant. More precisely, we take  $Y$  to be the space of continuous functions on the interval  $[0, 1]$  vanishing at the endpoints, consider a general class of linear measurements, and assume that  $\varepsilon$  is Gaussian white noise. Our choice of  $Y_n \subset Y$  is the space of piecewise linear continuous functions  $u_n$  specified by the point values

$$u_j^n := u_n(x_j^n), \quad x_j^n = \frac{j}{n+1} \quad \text{for } j = 0, \dots, n+1.$$

Further, we take  $\pi_{\text{pr}}^{(n)}$  in (1.3) to be the discrete TV prior distribution

$$\pi_{\text{pr}}^{(n)}(u_1^n, \dots, u_n^n) = c_n' \exp \left( -\alpha_n \sum_{j=1}^{n+1} |u_j^n - u_{j-1}^n| \right),$$

where  $\alpha_n > 0$  is a parameter and  $c_n' > 0$  is a normalization constant. This leads to the following posterior distribution:

$$\pi(u_1^n, \dots, u_n^n | \hat{m}) = c_n \exp \left( -\alpha_n \sum_{j=1}^{n+1} |u_j^n - u_{j-1}^n| - \frac{1}{2\sigma^2} \|Au_n - \hat{m}\|_{\mathbb{R}^N}^2 \right), \quad (1.4)$$

where  $c_n > 0$  is again a normalization constant.

Our main theorems 4.1 and 5.2 concern the behaviour of MAP and CM estimates for (1.4) when  $n \rightarrow \infty$ . Their proofs are based on epiconvergence of optimization problems and specific types of stochastic convergence, respectively. According to our theorems, there are essentially two choices of the parameter  $\alpha_n$  as a function of  $n$  leading to nontrivial convergence behaviour as  $n \rightarrow \infty$ : either  $\alpha_n = 1$ , or  $\alpha_n = \sqrt{n+1}$ . Let us summarize our results:

- If  $\alpha_n = 1$ , then the functions  $u_n^{\text{MAP}}$  converge in bounded variation (BV) topology. This case is equivalent to TV regularization. However, the TV prior distributions diverge, and so do the functions  $u_n^{\text{CM}}$ .
- If  $\alpha_n = \sqrt{n+1}$ , then the posterior distribution (1.4) converges to a distribution of a random variable  $v$  taking values in  $Y$ . However, condition (ii) above is not satisfied. Moreover,  $v$  is Gaussian<sup>3</sup>, and consequently  $u_n^{\text{CM}}$  is not edge-preserving with large  $n$ . The functions  $u_n^{\text{MAP}}$  converge to zero.
- If  $\alpha_n = (n+1)^r$  for any  $r \in \mathbb{R} \setminus \{0, \frac{1}{2}\}$ , then the TV prior distributions diverge and the functions  $u_n^{\text{MAP}}$  either converge to a useless limit or diverge.

We conclude that TV prior is not discretization invariant in the sense of (i) and (ii) above. Thus the answer to the question in the title is negative. However, we view our results positively as a quest for researchers to design discretization invariant prior distributions for edge-preserving inversion of (1.1).

What is the relevance of our results to practical Bayesian inversion? First, even if ‘blocky’ CM estimates are achieved by using the TV prior in a fixed discrete model space, refining the discretization (while keeping the measurement fixed) leads to smooth, non-blocky CM estimates. Second, inversion algorithms are often tuned and debugged using a coarse discretization before refining the model space for more accurate inversion. Expressing *a priori* information in a discretization invariant manner ensures that the results using coarse and fine discretization are consistent.

This paper is organized as follows. In section 2 we give basic definitions and show how regularization theory and Bayesian inversion are related. In section 3 we define discretization invariance. In section 4 we prove a result about convergence of MAP estimates. In section 5 we prove results about convergence of prior distributions and CM estimates. In section 6 we illustrate our theoretical findings by numerical computations.

## 2. The generic posterior distribution

We restrict ourselves in this work to a class of one-dimensional inverse problems. Our choice of spaces  $Y$  and  $Y_n$  is as follows.

**Definition 2.1** (function spaces  $Y$  and  $Y_n$ ). *Let  $Y$  be the space of continuous functions on the interval  $[0, 1]$  vanishing at the endpoints:*

$$Y = C_0([0, 1]) = \{u \in C([0, 1]) : u(0) = u(1) = 0\}.$$

*For any integer  $n > 0$ , let  $Y_n \subset Y$  be the following set of piecewise linear functions on the interval  $[0, 1]$ :*

$$Y_n = \{u \in Y : u|_{[x_j^n, x_{j+1}^n]} \text{ is linear for } j = 0, \dots, n\},$$

where

$$x_j^n = \frac{j}{n+1} \quad \text{for } j = 0, \dots, n+1.$$

<sup>3</sup> Numerical evidence and a conjecture was first presented by Markku Lehtinen [21].

Further, consider the roof-top basis  $\{\psi_j^n\}_{j=1}^n$  for  $Y_n$ , where  $\psi_j^n \in Y_n$  satisfy  $\psi_j^n(x_k^n) = \delta_{jk}$  for  $j = 1, \dots, n$  and  $k = 0, \dots, n + 1$ .

We use the following class of probability distributions as priors.

**Definition 2.2** (*p*-variation distribution). *Let  $(\Omega, \Sigma, P)$  be a complete probability space. Let  $n > 0$  be an integer,  $\alpha_n > 0$ , and  $1 \leq p \leq 2$ . The  $Y_n$ -valued random function*

$$u_n(t) = u_n(t, \omega) = \sum_{j=1}^n u_j^n(\omega) \psi_j^n(t), \quad \omega \in \Omega, \tag{2.1}$$

has *p*-variation distribution in  $Y_n$  if the  $\mathbb{R}^n$ -valued random vector  $[u_1^n(\omega), \dots, u_n^n(\omega)]^T$  has the probability density function

$$\begin{aligned} \pi_{p,n}(u_1^n, \dots, u_n^n) &= c_{p,n} \exp \left( -\alpha_n \left\| \frac{\partial}{\partial t} \sum_{j=1}^n u_j^n \psi_j^n(t) \right\|_{L^p(0,1)}^p \right) \\ &= c_{p,n} \exp \left( -\frac{\alpha_n}{(n+1)^{1-p}} \sum_{j=1}^{n+1} |u_j^n - u_{j-1}^n|^p \right), \end{aligned} \tag{2.2}$$

where  $\alpha_n > 0$  is a parameter,  $u_0^n = u_{n+1}^n = 0$  and  $c_{p,n}$  is a normalization constant. The special case  $p = 1$  is called the total variation (TV) probability distribution in  $Y_n$ .

Note that  $p = 2$  gives a Gaussian distribution. In definition 2.2 we require  $u_0^n = u_{n+1}^n = 0$  for the *p*-variation distribution to be a probability density function. Without this requirement, a constant could be added to  $u_n$  without altering  $\pi_{p,n}(u_1^n, \dots, u_n^n)$ .

We are ready to define the posterior distribution analysed in this paper.

**Definition 2.3** (the generic posterior distribution). *Denote by  $\mathcal{Z}$  the set of Borel measures  $A(dt)$  with finite variation that are supported on compact subsets of  $(0, 1)$ . Let  $A_j \in \mathcal{Z}$  for  $j = 1, \dots, N$  and  $\sigma > 0$ . Given  $u \in Y$ , let the measurement  $m$  be a random vector in  $\mathbb{R}^N$  with components*

$$m_j = (Au)_j + \varepsilon_j = \langle A_j, u \rangle + \varepsilon_j = \int_0^1 v(t) A_j(dt) + \varepsilon_j, \tag{2.3}$$

where  $\varepsilon_j \sim N(0, \sigma)$  are independent Gaussian errors. Assume as given a realization  $\hat{m}$  of  $m$ . Modelling a priori knowledge about  $u$  with the *p*-variation distribution in the space  $Y_n$  leads to the posterior distribution

$$\pi(u_n | \hat{m}) = \tilde{c} \exp \left( -\frac{1}{2\sigma^2} \|Au_n - \hat{m}\|_{\mathbb{R}^N}^2 - \alpha_n \|u_n'\|_p^p \right), \tag{2.4}$$

where  $\|\cdot\|_{\mathbb{R}^N}$  is the standard Euclidean norm and  $\|u_n'\|_p^p$  is understood in the sense of (2.1) and (2.2).

Recall that  $A(dt)$  has finite variation if there are finite non-negative measures  $A_1(dt)$  and  $A_2(dt)$  such that  $A(dt) = A_1(dt) - A_2(dt)$ . The norm of  $\mathcal{Z}$  is given by

$$\|A\|_{\mathcal{Z}} = \inf\{A_1([0, 1]) + A_2([0, 1]) : A(dt) = A_1(dt) - A_2(dt)\}.$$

We close the section by pointing out a connection between regularization theory and Bayesian MAP estimates. Maximizing the posterior distribution (2.4) is equivalent to the minimization problem

$$\arg \min_{u_n \in Y_n} \left\{ \|Au_n - m\|_{\mathbb{R}^N}^2 + \beta_n \int_0^1 |u_n'(t)|^p dt \right\} \tag{2.5}$$

with  $\beta_n = 2\sigma^2\alpha_n$ . But (2.5) is by definition the Tikhonov ( $p = 2$ ) or TV ( $p = 1$ ) regularized solution of the inverse problem (1.2). We note that the edge-preserving property of discrete TV regularization is related to the regularization term in (2.5) allowing large values of derivatives.

### 3. Discretization invariance

It is tempting to consider the  $p$ -variation prior distribution (2.2) as discretization of the following formal prior distribution:

$$\pi_u(v) \stackrel{\text{formally}}{\sim} \exp(-\alpha\|v'\|_p^p), \quad v \in Y. \quad (3.1)$$

Generalizing the successful solution of finite-dimensional inverse problems in  $Y_n$  to solution of the continuous problem in  $Y$  using (3.1) is a natural idea. This can indeed be done for  $p = 2$  (see Lehtinen *et al* [22] and Lasanen [20]). To analyse the case  $1 \leq p < 2$  we need the following definitions.

**Definition 3.1** (convergence i.d.). *Let  $X_n$  and  $X$  be random variables having probability density functions  $\pi_{X_n}$  and  $\pi_X$ , respectively. We say that  $X_n$  converges to  $X$  in distribution, or i.d., if  $\pi_{X_n}(x) \rightarrow \pi_X(x)$  as  $n \rightarrow \infty$  at every continuity point  $x$  of  $\pi_X(x)$ .*

**Definition 3.2** (linear discretization of random functions). *Let  $u_{n(\ell)}(t, \omega)$  be  $Y_{n(\ell)}$ -valued random variables with  $\ell = 1, 2, 3, \dots$  and  $1 < n(1) < n(2) < n(3) < \dots$ . Assume that*

(i) *There is a  $Y$ -valued random variable  $v(t) = v(t, \omega)$  such that for any  $t \in [0, 1]$*

$$\lim_{\ell \rightarrow \infty} u_{n(\ell)}(t) = v(t) \quad \text{i.d.}$$

(ii) *There are bounded linear operators  $T_{n(\ell)} \in L(Y)$  such that for any  $t \in [0, 1]$*

$$u_{n(\ell)}(t) = (T_{n(\ell)}v)(t).$$

*Then  $u_{n(\ell)}$  are linear discretizations of a random function. Further, we say that  $v$  can be approximated by finite-dimensional random variables in a discretization invariant manner and  $u_{n(\ell)}$  are proper linear discretizations of  $v$ .*

Note that definition 3.2 is analogous to that of Lasanen [20] (see also [15, 23]).

**Definition 3.3** (discretization invariant choice of  $Y_n$  and  $\pi_{\text{pr}}^{(n)}$ ). *Assume as given  $Y_{n(\ell)}$  and  $\pi_{\text{pr}}^{(n(\ell))}$  for (1.3) with  $\ell = 1, 2, 3, \dots$  and  $1 < n(1) < n(2) < n(3) < \dots$ . Let  $u_{n(\ell)}$  be random functions taking values in  $Y_{n(\ell)}$  and having distribution  $\pi_{\text{pr}}^{(n(\ell))}$ . We say that the choice of  $Y_{n(\ell)}$  and  $\pi_{\text{pr}}^{(n(\ell))}$  is discretization invariant if  $u_{n(\ell)}$  are linear discretizations of a random function in the sense of definition 3.2.*

We will show that the discrete random variables distributed according to the generic posterior distribution (2.4) with  $1 \leq p < 2$  are not linear discretizations of any random function. See remark 5.1 on page 12.

#### 4. Convergence of MAP estimates

We analyse the convergence of MAP estimates

$$u_n^{\text{MAP}}(t; p, \alpha_n) \in \arg \max_{u_n \in Y_n} \exp \left( -\frac{1}{2\sigma^2} \|Au_n - \hat{m}\|_{\mathbb{R}^N}^2 - \alpha_n \|u_n'\|_p^p \right) \quad (4.1)$$

for the posterior distribution (2.4) as the discretization is refined arbitrarily:  $n = n(\ell) = 2^\ell - 1$  and  $\ell \rightarrow \infty$ . This choice of  $n(\ell)$  ensures that  $Y_{n(\ell)} \subset Y_{n(\ell+1)}$ , which is needed below in the case  $p = 1$ . In the case  $1 < p \leq 2$  the object function in (4.1) is strictly convex and there is a unique MAP estimate, whereas in the case  $p = 1$  the MAP estimate is not necessarily unique.

We recall two important function spaces. First, the Sobolev space  $W^{1,p}(0, 1)$  consists of  $L^p(0, 1)$  functions with weak derivatives in  $L^p(0, 1)$ . By the Sobolev imbedding theorem we know that  $W_0^{1,p}$  functions are continuous. Second, the space of functions of bounded variation is defined as follows. Let

$$\begin{aligned} BV(0, 1) &= \{u \in L^1(\mathbb{R}) : \|u\|_{BV} < \infty, \text{supp}(u) \subset [0, 1]\}, \\ BV_0(0, 1) &= \{u \in BV(0, 1) : u(0) = u(1) = 0\}, \end{aligned}$$

where

$$\|u\|_{BV} = \sup \left\{ \int_0^1 u(s) \partial_s \phi(s) \, ds : \phi \in C^\infty([0, 1]), \|\phi\|_{L^\infty} \leq 1 \right\}.$$

Note that for  $u \in BV_0(0, 1)$  the derivative  $\partial_s u$ , defined in the sense of distributions, is a measure. We say that  $u_n \in BV(0, 1)$  converge in weak\* topology of  $BV$  to  $u \in BV(0, 1)$  if  $\|u_n - u\|_{L^1} \rightarrow 0$  and  $\int_{\mathbb{R}} \phi(\partial_s u_n - \partial_s u)(s) \rightarrow 0$  for any  $\phi \in C_0([0, 1])$ , i.e., the measures  $\partial_s u_n$  converge weakly to  $\partial_s u$ . In this case we denote  $u_n \xrightarrow{BV-w^*} u$ . Note that the trace  $u \mapsto u(s_0)$ ,  $0 \leq s_0 \leq 1$  is continuous from weak\* topology of  $BV(0, 1)$  to  $\mathbb{R}$ .

**Theorem 4.1.** *Let  $\sigma > 0$ ,  $\hat{m} \in \mathbb{R}^N$  and  $n = n(\ell) = 2^\ell - 1$  for  $\ell = 2, 3, \dots$ . Assume  $A_j \in L^1(0, 1) \cap \mathcal{Z}$  for  $j = 1, \dots, N$ .*

(i) *Let  $\alpha_n = \tilde{\alpha}/(2\sigma^2)$  with some  $\tilde{\alpha} > 0$  and  $1 < p \leq 2$ . Let  $u_n^{\text{MAP}} = u_n^{\text{MAP}}(t; p, \alpha_n)$  be the unique MAP estimate given by (4.1) for the posterior distribution (2.4). Then there is a unique limit function  $\tilde{u}(\cdot; p, \tilde{\alpha}) \in W_0^{1,p}(0, 1)$  such that*

$$\lim_{\ell \rightarrow \infty} u_{n(\ell)}^{\text{MAP}} = \tilde{u}(\cdot; p, \tilde{\alpha})$$

*in the weak topology of  $W_0^{1,p}(0, 1)$ .*

(ii) *Let  $\alpha_n = \tilde{\alpha}/(2\sigma^2)$  with some  $\tilde{\alpha} > 0$  and  $p = 1$ . Then for any sequence  $\{u_{n(\ell)}^{\text{MAP}}\}_{\ell=2}^\infty$  of maximizers of (4.1) there is a subsequence that converges in weak\* topology of  $BV$  to some  $\tilde{u} \in BV_0(0, 1)$ .*

(iii) *Let the parameters  $\alpha_n > 0$  satisfy  $\lim_{\ell \rightarrow \infty} \alpha_{n(\ell)} = \infty$ . Then*

$$\lim_{\ell \rightarrow \infty} u_{n(\ell)}^{\text{MAP}} = 0$$

*in the norm topology of  $W_0^{1,p}(0, 1)$  for  $1 < p \leq 2$  and  $BV$  for  $p = 1$ .*

We note that the restriction  $A_j \in L^1(0, 1)$  is needed only in the case  $p = 1$ . The proof of theorem 4.1 consists in part of standard arguments in the field of epiconvergence of minimization problems. However, we present the details for the reader's convenience.

**Proof.** We define two optimization problems that are limits of optimization problems in finite-dimensional spaces. In the case  $1 < p \leq 2$  we consider the problem

$$\tilde{u}(t; p, \beta) = \arg \min_{u \in W_0^{1,p}} S(u), \quad S(u) = \|Au - \hat{m}\|_{\mathbb{R}^N}^2 + \beta \|u'\|_p^p, \quad (4.2)$$

where  $\beta > 0$  and  $W_0^{1,p}$  is the closure of  $C_0^\infty(0, 1)$  in the  $W^{1,p}(0, 1)$  norm. In the case  $p = 1$  we consider the problem

$$\tilde{u}(t; p, \beta) \in \arg \min_{u \in BV_0(0,1)} S(u), \quad S(u) = \|Au - \hat{m}\|_{\mathbb{R}^N}^2 + \beta \|u\|_{BV}. \quad (4.3)$$

Next we use methods of convex analysis and approximate (4.2), (4.3) with a discrete minimization problem

$$\tilde{u}_n(t; p, \beta_n) \in \arg \min_u S_n(u), \quad (4.4)$$

where  $S_n : W_0^{1,p} \rightarrow \overline{\mathbb{R}}_+ = \mathbb{R}_+ \cup \{0, \infty\}$  for  $1 < p \leq 2$  or  $S_n : BV_0(0, 1) \rightarrow \overline{\mathbb{R}}_+$  for  $p = 1$  is the convex nonlinear function

$$S_n(u) = \|Au - \hat{m}\|_{\mathbb{R}^N}^2 + \beta_n \|u'\|_p^p + I_{Y_n}(u).$$

Here  $\beta_n > 0$  are such that  $\beta = \lim_{n \rightarrow \infty} \beta_n$  and  $I_{Y_n}$  is the convex indicator function:  $I_{Y_n}(u) = 0$  if  $u \in Y_n$  and  $I_{Y_n}(u) = \infty$  if  $u \notin Y_n$ .

The proof of the convergence of the MAP estimate for the posterior distribution (2.4) is based, using the terminology of [28], to the epiconvergence of  $S_n$  to  $S$ .

Note that (4.1) is equivalent to

$$u_n^{\text{MAP}}(t; p, \alpha_n) = \arg \min_{u_n \in Y_n} (\|Au_n - \hat{m}\|_{\mathbb{R}^N}^2 + 2\sigma^2 \alpha_n \|u_n'\|_p^p), \quad (4.5)$$

which in turn is equivalent to (4.4) with  $\beta_n = 2\sigma^2 \alpha_n = \tilde{\alpha}$ .

Consider the operators  $T_n : C_0(0, 1) \rightarrow Y_n$  defined by

$$T_n u(t) = \sum_{j=1}^n u(x_j^n) \psi_j^n(t), \quad x_j^n = \frac{j}{n+1},$$

where the functions  $\psi_j^n$  are as in definition 2.1. Thus  $T_n u$  is obtained by linear interpolation from the point values  $u(x_j^n)$ .

*Properties of  $T_n$  for  $1 < p \leq 2$ .* Denote  $\Delta x^n := x_2^n - x_1^n = (n+1)^{-1}$ . Now for  $t \in I = I_j^n = [x_j^n, x_{j+1}^n]$  with  $j = 0, 1, 2, \dots, n$ , we have

$$T_n u(t) = u(x_j^n) + \frac{t - x_j^n}{\Delta x^n} \int_{x_j^n}^{x_{j+1}^n} u'(t) dt,$$

and, further,

$$(T_n u)'(t) = \frac{1}{\Delta x^n} \int_{x_j^n}^{x_{j+1}^n} u'(t) dt = [u']_I,$$

where we denote by  $[u']_I$  the average of  $u'$  over  $I$ . Note that  $|(T_n u)'(t)| \leq 2Mu'(t)$  where  $Mu'$  is the Hardy–Littlewood maximal function of  $u'$ . For any  $t \in (0, 1)$ , let  $j(t, n)$  be some index for which  $t \in I_{j(t,n)}^n$ . Since  $u' \in L^1(0, 1)$ , we have (similarly to the standard theorem of Lebesgue points)

$$\lim_{n \rightarrow \infty} u'(t) - [u']_{I_{j(t,n)}^n} = 0$$

for almost every  $t$ , see [10]. Since  $|(T_n u)'(t) - u'(t)| \leq (2Mu'(t) + |u'(t)|) \in L^p(0, 1)$ , Lebesgue's theorem of dominated convergence yields that

$$\lim_{n \rightarrow \infty} \|(T_n u - u)'\|_{L^p(0,1)}^p = \int_0^1 \lim_{n \rightarrow \infty} |u'(t) - (u')_{I_{j(t,n)}^n}|^p dt = 0.$$

Hence  $\lim_{n \rightarrow \infty} T_n = I$  in the strong operator topology of  $W_0^{1,p}$ . Moreover,

$$\|(T_n u)'\|_{L^p} \leq 2\|Mu'\|_{L^p} \leq C\|u'\|_{L^p}$$

and hence the norms  $\|T_n\|_{W_0^{1,p} \rightarrow W_0^{1,p}}$  are uniformly bounded.

*Epicomvergence in the case  $1 < p \leq 2$ .* First we consider the case  $\beta_n = \beta = \tilde{\alpha}$ . Let  $u_n = u_{n(\ell)}$ ,  $\ell = 2, 3, \dots$ , be a converging sequence in  $W_0^{1,p}$  with  $\lim_{n \rightarrow \infty} u_n = u$ . We have shown that  $\lim_{n \rightarrow \infty} T_n = I$  in the strong operator topology of  $W_0^{1,p}$  and that the norms  $\|T_n\|_{W_0^{1,p} \rightarrow W_0^{1,p}}$  are uniformly bounded. Thus  $\lim_{n \rightarrow \infty} T_n u_n = u$ . Further, since  $S_n$  has infinite values in the complement of  $Y_n$  and  $T_n|_{Y_n} = I$ , we see trivially that  $S_n(u_n) \geq S_n(T_n u_n)$ . Using these facts we estimate

$$\begin{aligned} \liminf_{n \rightarrow \infty} S_n(u_n) &\geq \liminf_{n \rightarrow \infty} S_n(T_n u_n) \\ &\geq \liminf_{n \rightarrow \infty} \|A(T_n u_n - u) + (Au - \hat{m})\|_{\mathbb{R}^N}^2 + \beta \|(T_n u_n)'\|_p^p \\ &\geq S(u). \end{aligned} \tag{4.6}$$

Moreover, for  $u \in W_0^{1,p}$  there is such a sequence  $v_n = T_n u \rightarrow u$  in  $W_0^{1,p}$  that

$$\limsup_{n \rightarrow \infty} S_n(v_n) = \limsup_{n \rightarrow \infty} \|A(T_n u_n - u) + (Au - \hat{m})\|_{\mathbb{R}^N}^2 + \beta \|(T_n u)'\|_p^p = S(u). \tag{4.7}$$

By definition (see [3], property 1.14 and also [2, 28]), formulae (4.6) and (4.7) mean that the functions  $S_n$  epiconverge to  $S$ . Define  $s_0 = \inf \{S(v) : v \in W_0^{1,p}\}$  and  $s_n = \inf \{S_n(v) : v \in W_0^{1,p}\}$  for  $n = n(\ell)$ ,  $\ell = 2, 3, 4, \dots$ , and consider the set

$$\operatorname{argmin}(S_n) = \{u \in W_0^{1,p} : S_n(u) = s_n\},$$

where  $S_n$  attains its minimum (actually, this set contains only the function  $\tilde{u}_n(\beta_n)$ ). Since  $S_n$  epiconverge to  $S$  [3], proposition 2.9 yields that

$$\limsup_{n \rightarrow \infty} \operatorname{argmin}(S_n) \subset B_\varepsilon = \{u \in W_0^{1,p} : S(u) \leq s_0 + \varepsilon\} \tag{4.8}$$

for any  $\varepsilon > 0$ . Now assume that  $0 < \beta < \infty$ . Since  $S$  is strictly convex and weakly lower semicontinuous,  $S$  has a unique global minimum at  $\tilde{u}$  (see e.g. [5, theorem 2.1.4]). Thus (4.8) implies that  $\lim_{n \rightarrow \infty} S(u_{n(\ell)}^{\text{MAP}}) = \min(S)$  and in particular  $\|u_{n(\ell)}^{\text{MAP}}\|_{W_0^{1,p}}$  are uniformly bounded.

Assume now that  $u_{n(\ell)}^{\text{MAP}}$  do not converge weakly to  $\tilde{u}$ . Then there is  $f \in (W_0^{1,p})'$  such that for some subsequence  $\lim_{k \rightarrow \infty} \langle f, u_{n(\ell_k)}^{\text{MAP}} \rangle = c_f \neq \langle f, \tilde{u} \rangle$ . By the Banach–Alaoglu theorem  $\{u_{n(\ell_k)}^{\text{MAP}}\}$  has a weakly converging subsequence. We denote the limit of such a subsequence by  $\tilde{u}_1$ . Since  $S : W_0^{1,p}(0, 1) \rightarrow \mathbb{R}$  is lower semicontinuous in the weak topology of  $W_0^{1,p}$ , we see that  $S(\tilde{u}_1) = \min(S)$ . Then  $\langle \tilde{u} - \tilde{u}_1, f \rangle \neq 0$  is in contradiction with the fact that the minimum  $\tilde{u}$  is unique. This shows that  $u_{n(\ell)}^{\text{MAP}}$  converges weakly in  $W_0^{1,p}(0, 1)$  to  $\tilde{u}$ . This proves (i).

*Approximation of functions in the case  $p = 1$ .* First we consider approximations in  $W_0^{1,1}(0, 1) \subset BV_0(0, 1)$ . Let  $h \in L^1(0, 1)$  and  $\mathcal{N}_\ell$  be the  $\sigma$ -algebra generated by intervals  $(0, x_j^{n(\ell)})$ ,  $j = 0, 1, \dots, n(\ell) + 1$ . Then the  $\sigma$ -algebra  $\mathcal{N}$  generated by  $\cup_{\ell=2}^\infty \mathcal{N}_\ell$  is the standard



Borel  $\sigma$ -algebra of  $(0, 1)$ . Since  $\mathcal{N}_{\ell+1} \subset \mathcal{N}_\ell$ , it follows from Doob’s second martingale theorem ([27], corollary C.9, [11], theorem 10.5.7) that

$$\lim_{\ell \rightarrow \infty} \|\mathbb{E}(h|\mathcal{N}_\ell) - h\|_{L^1(0,1)} = 0, \tag{4.9}$$

where  $\mathbb{E}(h|\mathcal{N}_\ell)$  is the conditional expectation with respect to the  $\sigma$ -algebra  $\mathcal{N}_\ell$ . Now, for  $u \in W_0^{1,1}(0, 1)$  the function  $(T_n u)'$  is piecewise constant and

$$(T_{n(\ell)} u)' = \mathbb{E}(u'|\mathcal{N}_\ell). \tag{4.10}$$

Formulae (4.9) and (4.10) imply

$$\lim_{\ell \rightarrow \infty} \|T_{n(\ell)} f - f\|_{W_0^{1,1}(0,1)} = 0. \tag{4.11}$$

Moreover, we see that  $\|T_n f\|_{W_0^{1,1}} = \|(T_n f)'\|_{L^1} \leq c\|f\|_{W_0^{1,1}}$ .

Second, we apply the properties of operators  $T_n$  to approximate functions in  $BV_0(0, 1)$ . By [36, theorem 5.2.1] for any  $u \in BV_0(0, 1)$  and any  $u_k \xrightarrow{BV-w^*} u$  we have

$$\|u\|_{BV} \leq \liminf_{k \rightarrow \infty} \|u_k\|_{BV}. \tag{4.12}$$

Let  $u \in BV_0(0, 1)$ . By [8, theorem 2.3], there are  $\Phi_k \in W^{1,1}(0, 1) \cap C^\infty((0, 1))$  such that  $\Phi_j$  converge to  $u$  in the weak\* topology of  $BV(0, 1)$  and

$$\lim_{k \rightarrow \infty} \|\Phi_k\|_{BV} = \|u\|_{BV}. \tag{4.13}$$

Note that  $\Phi_j$  are not assumed to vanish at boundary points. However, since the trace  $\phi \mapsto (\phi(0), \phi(1))$  is weakly continuous map  $BV(0, 1) \rightarrow \mathbb{R}^2$  we see that  $\phi_j(s) = \Phi_j(s) - (1 - s)\Phi_j(0) - s\Phi_j(1) \in W_0^{1,p}(0, 1)$  converge to  $u$  in the weak\* topology of  $BV_0(0, 1)$ . Moreover, by (4.13)

$$\lim_{j \rightarrow \infty} \|\phi_j\|_{BV} = \|u\|_{BV}. \tag{4.14}$$

By (4.11), there are  $\ell_j, \ell_{j+1} > \ell_j$  such that for any  $j$  and  $\ell \geq \ell_j$  we have

$$\|T_{n(\ell)} \phi_j - \phi_j\|_{W_0^{1,1}(0,1)} \leq \frac{1}{j}.$$

This implies that there the sequence  $T_{n(\ell_j)} \phi_j$  converges to  $u$  in the weak\* topology of  $BV$  and

$$\lim_{j \rightarrow \infty} \|T_{n(\ell_j)} \phi_j\|_{BV} = \|u\|_{BV}.$$

In the following, we denote  $u_{n_j} = T_{n_j} \phi_j, n_j = n(\ell_j)$ .

*Epiconvergence in the case  $p = 1$ .* Since  $A_k \in L^1$  and by [5, p 41], the embedding  $BV_0 \rightarrow L^\infty(0, 1)$  from the weak\* topology of  $BV$  to the norm topology of  $L^\infty$  is continuous, we see that  $\langle A_k, u_{n_j} \rangle \rightarrow \langle A_k, u \rangle$ . Thus, for any  $u \in BV_0(0, 1)$  there are  $u_{n_j} \in Y_{n_j}$  such that

$$S(u) = \limsup_{j \rightarrow \infty} S_{n_j}(u_{n_j}). \tag{4.15}$$

Next we note that  $\lim_{k \rightarrow \infty} \langle A_j, u_k \rangle \rightarrow \langle A_j, u \rangle$ . Combining this with (4.12) we see that for any  $u \in BV_0(0, 1)$  and any  $u_k \xrightarrow{BV-w^*} u$  we have

$$S(u) \leq \liminf_{k \rightarrow \infty} S_k(u_k). \tag{4.16}$$

Again, (4.15), (4.16) imply that  $S_n$  epiconverge to  $S$ . Thus by [3], proposition 2.9,

$$\limsup_{n \rightarrow \infty} \operatorname{argmin}(S_n) \subset B_\varepsilon = \{u \in BV_0 : S(u) \leq s_0 + \varepsilon\}. \tag{4.17}$$

Thus if  $u_k = u_{n(k)}^{\text{MAP}} \in \operatorname{argmin}(S_{n(k)})$ , we see that  $\lim_{k \rightarrow \infty} S(u_k) = \inf(S)$ . Moreover,  $\|u_k\|_{BV}$  are uniformly bounded. The sequence  $u_k$  has a subsequence that converges weakly (see

[5, p 41]). For more detail, for proof of  $L^1$  convergence, see [36, corollary 5.3.4] and for the fact that the measures converge weakly, use the Riesz representation theory and the Banach–Alaoglu theorem. If  $\tilde{u}$  is a limit of such a subsequence we have  $S(\tilde{u}) = \inf(S)$ . By (4.16) for any converging subsequence the limit is a minimizer of  $S$ . This proves (ii).

Finally, we consider (iii). We see that if  $\lim_{n \rightarrow \infty} \beta_n = \infty$  then  $\lim_{n \rightarrow \infty} \|u_n^{\text{MAP}}\|_{W_0^{1,p}} = 0$  for  $1 < p \leq 2$  and  $\lim_{n \rightarrow \infty} \|u_n^{\text{MAP}}\|_{BV} = 0$  for  $p = 1$  proving the assertion.  $\square$

### 5. Convergence of the CM estimate

We analyse the convergence of the posterior distribution (2.4) depending on parameter  $\alpha_n$  as the discretization is refined arbitrarily, or  $n \rightarrow \infty$ . In particular, we are interested in the convergence of the CM estimate

$$u_n^{\text{CM}}(t; p, \alpha_n) = \int_{\mathbb{R}^n} \left( \sum_{j=1}^n s_j \psi_j^n(t) \right) \pi(s_1, \dots, s_n | \hat{m}) ds_1 \cdots ds_n, \tag{5.1}$$

where  $\pi(s_1, \dots, s_n | \hat{m})$  is the conditional probability density function of coefficients of  $u_n$  in the basis  $\{\psi_j^n\}$ .

We introduce some definitions and notation. In the following we adapt the standard notation by sometimes omitting the variable  $\omega$ , i.e., we write  $v(t, \omega) = v(t)$  where  $t \in [0, 1]$  and  $\omega \in \Omega$ .

**Definition 5.1** (convergence weakly i.d.). *Let  $v_n$  and  $v$  be  $C(0, 1)$ -valued random variables. We say that  $v_n$  converges to  $v$  weakly in distribution if  $\langle A, v_n \rangle \rightarrow \langle A, v \rangle$  i.d. for all  $A \in \mathcal{Z}$  when  $n \rightarrow \infty$ .*

**Definition 5.2** (convergence i.p.). *Let  $v_n$  and  $v$  be  $C(0, 1)$ -valued random variables. We say that  $v_n$  converges to  $v$  in probability, or i.p., if for every  $\varepsilon > 0$  we have  $P\{\|v_n - v\| > \varepsilon\} \rightarrow 0$  as  $n \rightarrow \infty$ .*

**Definition 5.3** (uniform integrability). *Let  $v_n$  be  $C(0, 1)$ -valued random variables for  $n = 1, 2, 3, \dots$ . We say that the family  $v_n$  is uniformly integrable if*

$$\sup_n \mathbb{E}(|v_n| 1_{\{|v_n|>c\}}) \rightarrow 0, \quad c \rightarrow \infty,$$

where  $1_{\{|v_n|>c\}}$  is the indicator function equal to 1 if  $|v_n| > c$  and zero otherwise.

**Definition 5.4** (measurement  $\sigma$ -algebras  $\mathcal{M}, \mathcal{M}_n$ ). *Denote  $m_j = \langle A_j, v \rangle + \varepsilon_j$  and  $m_j^n = \langle A_j, v_n \rangle + \varepsilon_j$  with some  $A_j \in \mathcal{Z}$  for  $j = 1, \dots, N$  and independent errors  $\varepsilon_j \sim N(0, 1)$ . Let the  $\sigma$ -algebras  $\mathcal{M}, \mathcal{M}_n \subset \Sigma$  be generated by the sets*

$$\begin{aligned} \{\omega \in \Omega : m_j(\omega) < \lambda_j\}, & \quad \lambda_j \in \mathbb{R}, \\ \{\omega \in \Omega : m_j^n(\omega) < \lambda_j\}, & \quad \lambda_j \in \mathbb{R}, \end{aligned}$$

respectively.

Note that the above noise processes  $\varepsilon_j$  are the same for  $m_j$  and  $m_j^n$ , and for simplicity we take  $\sigma = 1$  in this section.

In the following, we use the abbreviations a.e. and a.s. for the terms *almost every* and *almost surely*, respectively.

We denote the conditional expectation of  $v(t)$  with respect to  $\sigma$ -algebra  $\mathcal{M}$  by  $\mathbb{E}(v(t)|\mathcal{M})$ . Recall that  $\mathbb{E}(v(t)|\mathcal{M}) = \mathbb{E}(v(t)|\mathcal{M})(\omega)$  is the random variable that is measurable with respect to  $\mathcal{M}$  and for which

$$\int_S \mathbb{E}(v(t)|\mathcal{M})P(d\omega) = \int_S v(t)P(d\omega) \quad \text{for all } S \in \mathcal{M}.$$

Since  $\mathbb{E}(v(t)|\mathcal{M})$  is a  $\mathcal{M}$ -measurable random variable, there exists a deterministic function  $\hat{m} \mapsto \tilde{E}(v(t)|\hat{m})$  so that

$$\mathbb{E}(v(t)|\mathcal{M}) = \tilde{E}(v(t)|m(\omega)) \quad \text{a.s.}$$

(see [11], theorem 4.2.8). We call  $\tilde{E}(v(t)|\hat{m})$  the conditional mean with measurement  $\hat{m}$  and occasionally denote it by  $\tilde{E}(v(t)|m = \hat{m})$ . Let  $B(\hat{m}, r) \subset \mathbb{R}^N$  be a ball with centre  $\hat{m}$  and radius  $r$ . We can write

$$\tilde{E}(v(t, \omega)|\hat{m}) = \lim_{r \rightarrow 0} \frac{1}{P(\{m(\omega) \in B(\hat{m}, r)\})} \int_{\{m(\omega) \in B(\hat{m}, r)\}} v(t, \omega)P(d\omega) \quad (5.2)$$

for a.e.  $\hat{m}$ . If a conditional probability density function  $\pi_{v(t)}(\cdot | \hat{m})$  exists and  $\varepsilon$  is normally distributed, the Radon–Nikodym derivative of the density of  $m$  with respect to Lebesgue measure is  $C^\infty$  smooth and we can write

$$\tilde{E}(v(t)|\hat{m}) = \int_{\mathbb{R}} s \pi_{v(t)}(s|\hat{m}) ds.$$

We define the conditional expectation  $\tilde{E}(v_n(t)|\hat{m})$  similarly.

**Theorem 5.1.** *Let  $v_n$  and  $v$  be random variables taking values in  $C(0, 1)$ . Assume that  $v_n \rightarrow v$  weakly i.d. when  $n \rightarrow \infty$  such that for a given  $t \in (0, 1)$  variables  $v_n(t)$ ,  $n = 1, 2, \dots$  are uniformly integrable and  $v(t, \omega)$ ,  $v_n(t, \omega) \in L^1(\Omega)$ . Then*

$$\lim_{n \rightarrow \infty} \tilde{E}(v_n(t)|\hat{m}) = \tilde{E}(v(t)|\hat{m}) \text{ for a.e. } \hat{m} \in \mathbb{R}^N, \quad (5.3)$$

$$\lim_{n \rightarrow \infty} \tilde{E}(F(v_n(t))|\hat{m}) = \tilde{E}(F(v(t))|\hat{m}) \text{ for a.e. } \hat{m} \in \mathbb{R}^N \quad (5.4)$$

where  $F = \chi_{(-\infty, \lambda]}$ ,  $\lambda \in \mathbb{R}$ .

The proof is given in section 5.2.

In particular, we consider the case where  $v_n = u_n$  are  $p$ -variation random variables taking values in  $Y_n$  and  $v = u_B$  is the Brownian bridge defined below.

**Definition 5.5** (Brownian bridge  $u_B$ ). *Define  $u_B$  as stochastic process  $u_B(t) = u_B(t, \omega)$ ,  $t \in [0, 1]$ ,  $\omega \in \Omega$ , having zero expectation and covariance function*

$$\mathbb{E}(u_B(t_1)u_B(t_2)) = \sigma_p^2 |t_1| |1 - t_2|, \quad (5.5)$$

where  $t_1, t_2 \in [0, 1]$ ,  $t_1 \leq t_2$  and

$$\sigma_p^2 = \int_{\mathbb{R}} x^2 e^{-|x|^p} dx. \quad (5.6)$$

By Kolmogorov's theorem and (5.5) we can choose such a version of  $u_B$  that its realizations  $t \mapsto u_B(t, \omega)$  are continuous a.s. Here we say that a random variable  $a(t, \omega)$  is a version of  $b(t, \omega)$  if the distributions of  $(a(t_1), \dots, a(t_\ell))$  and  $(b(t_1), \dots, b(t_\ell))$  in  $\mathbb{R}^\ell$  coincide for any  $t_1, \dots, t_\ell \in [0, 1]$  and  $\ell > 0$ .

**Theorem 5.2.** Let  $1 \leq p \leq 2$  and  $\tilde{\alpha} > 0$ . Let  $u_n, n = 1, 2, \dots$  be  $p$ -variation random functions in  $Y_n$  with parameter

$$\alpha_n = \tilde{\alpha}(n + 1)^{1-\frac{p}{2}}.$$

Then for any  $t \in [0, 1]$  the prior distributions converge, i.e.,

$$\lim_{n \rightarrow \infty} u_n(t) = u_B(t) \quad \text{i.d.} \tag{5.7}$$

Moreover, the posterior distributions converge in distribution:

$$\lim_{n \rightarrow \infty} \tilde{E}(F(u_n(t))|\hat{m}) = \tilde{E}(F(u_B(t))|\hat{m}) \quad \text{for a.e. } \hat{m} \in \mathbb{R}^N,$$

where  $F = \chi_{(-\infty, \lambda]}$ ,  $\lambda \in \mathbb{R}$  and the CM-estimates converge:

$$\lim_{n \rightarrow \infty} \tilde{E}(u_n(t)|\hat{m}) = \tilde{E}(u_B(t)|\hat{m}) \quad \text{for a.e. } \hat{m} \in \mathbb{R}^N.$$

Further, consider the parameter  $\alpha_n = \tilde{\alpha}(n + 1)^q$ . If  $q > 1 - \frac{p}{2}$  then  $\lim_{n \rightarrow \infty} u_n(t) = 0$ . If  $q < 1 - \frac{p}{2}$  then the random variables  $u_n(t)$  do not converge even in distribution.

The proof is given in section 5.2.

**Remark 5.1.** In (5.7) the limit  $u_B(t)$  is independent of  $p$  (up to the scaling factor  $\sigma_p$ ). Thus the choice  $p = 1$  represents in the limit  $n \rightarrow \infty$  the same *a priori* knowledge than the choice  $p = 2$  (Gaussian smoothness prior). In particular, this implies that the TV prior distribution is not discretization invariant: since TV prior distributions converge to a Gaussian distribution and any linear discretizations of Gaussian distributions are also Gaussian, we see that TV priors are not discretizations of any random variable in the sense of definition 3.2.

5.1. Convergence of TV prior distributions

Here we consider random variables  $u_n$  having  $p$ -variation distribution in  $Y_n$  and show that they converge to the Brownian bridge. This is needed to show that theorem 5.1 implies theorem 5.2.

We note that such results are well known in statistical mechanics—indeed, a non-harmonic random field in a one-dimensional lattice (such as  $u_n$ ) is generally known to converge to a free Gaussian field. For this type of result, see [7, 26]. However, compared to such work, we assume less regularity of the probability density functions and consider the integrals of these variables. For these reasons we think it is appropriate to present a full proof.

**Theorem 5.3.** Let  $1 \leq p \leq 2$  and  $\tilde{\alpha} > 0$ . Let  $u_n$  be a random variable taking values in  $Y_n$  and having  $p$ -variation distribution with parameter  $\alpha_n = \tilde{\alpha}(n + 1)^{1-\frac{p}{2}}$ . Moreover, let  $0 < t_1 < t_2 < \dots < t_Q < 1$ . Then

(i) We have

$$\lim_{n \rightarrow \infty} (u_n(t_1), \dots, u_n(t_Q)) = (U_{t_1}, \dots, U_{t_Q}) \quad \text{i.d.}$$

Here  $U_t$  are Gaussian random variables having Gaussian joint distributions with zero expectation and covariances

$$\mathbb{E}(U_{t_j}U_{t_k}) = \sigma_p^2|t_j||1 - t_k| \quad \text{for } t_j \leq t_k,$$

where  $\sigma_p$  is given by (5.6).

(ii) The variables  $u_n(t_j), j = 1, \dots, Q, n = 1, 2, \dots$ , are uniformly integrable.

(iii) Let  $0 < s' < t_1 < \dots < t_Q \leq s'' < 1$ . Let  $A_j \in \mathcal{Z}, j = 1, 2, \dots, N$  be measures supported on  $[s', s''] \subset (0, 1)$  and denote

$$a_n^j(\omega) = \int_{s'}^{s''} u_n(t, \omega) A_j(dt), \quad a^j(\omega) = \int_{s'}^{s''} u_B(t, \omega) A_j(dt).$$

Then  $a_n^j$  converges to  $a^j$  i.d. when  $n \rightarrow \infty$  for  $j = 1, \dots, N$ . Also, joint distributions of  $a_n^j$  and  $u_n(t_k)$  converge i.d. to  $a^j$  and  $u_B(t_k)$ , respectively, for  $j = 1, \dots, N$  and  $k = 1, \dots, Q$ . Here  $u_B$  is the Brownian bridge of definition 5.5.

**Proof.** Let  $(h_1^n, h_2^n, \dots, h_{n+1}^n)$  be a random vector in  $\mathbb{R}^{n+1}$  with density

$$\tilde{\pi}_n(y_1, y_2, \dots, y_{n+1}) = c \exp \left\{ -\tilde{\alpha}(n+1)^{p/2} \left( |y_1|^p + \sum_{j=1}^n |y_{j+1} - y_j|^p \right) \right\}. \tag{5.8}$$

Comparison of (5.8) and (2.2) shows that  $\tilde{\pi}_n(y_1, \dots, y_n, 0) = c_n \pi_{p,n}(y_1, \dots, y_n)$ . Define a piecewise linear function

$$h_n(t) = \sum_{j=1}^{n+1} h_j^n \psi_j^n(t), \quad t \in [0, 1], \tag{5.9}$$

where  $\psi_j^n \in Y^n$  are as in definition 2.1 for  $j = 1, \dots, n$ , and  $\psi_{n+1}^n$  is the piecewise linear function satisfying  $\psi_{n+1}^n(1) = 1$  and  $\psi_{n+1}^n(t) \equiv 0$  for  $0 \leq t \leq n/(n+1)$ .

Consider the convergence of a single variable  $u_n(t)$  for fixed  $0 < t < 1$ . Let

$$\xi_j^n = \tilde{\alpha}^{1/p}(n+1)^{1/2} (h_j^n - h_{j-1}^n), \tag{5.10}$$

where  $h_0^n = 0$  and  $j = 1, \dots, n+1$ . Then

$$h_j^n = \frac{1}{\tilde{\alpha}^{1/p}(n+1)^{1/2}} \sum_{\ell=1}^j \xi_\ell^n \quad \text{for } j = 1, \dots, n+1. \tag{5.11}$$

Now  $\xi_j^n$  are identically distributed independent variables with

$$\xi_j^n \sim \pi_\xi(t) := c_p \exp(-|t|^p), \quad j = 1, \dots, n+1. \tag{5.12}$$

Let  $\xi_\ell \sim \pi_\xi$  be independent variables for  $\ell = 1, 2, 3, \dots$  and define

$$S_j = \frac{1}{\sqrt{j}} \sum_{\ell=1}^j \xi_\ell \quad \text{for } j = 1, 2, 3, \dots$$

Note that  $S_j$  does not depend on  $n$ . Then  $h_n$  can be represented as

$$h_n(t) = k_{n,t} S_{\theta(n,t)} + r_{n,t} \xi_{\theta(n,t)+1}^n, \tag{5.13}$$

where

$$k_{n,t} = \frac{\theta(n,t)^{1/2}}{\tilde{\alpha}^{1/p}(n+1)^{1/2}}, \quad r_{n,t} = \frac{(n+1)t - \theta(n,t)}{\tilde{\alpha}^{1/p}(n+1)^{1/2}},$$

and  $\theta(n,t)$  is the largest integer  $j$  such that  $\frac{j}{n+1} \leq t$ .

For clarity, we (somewhat non-standardly) denote the probability density function of  $h_n(t)$  at  $a \in \mathbb{R}$  with the condition  $g = 0$  by

$$\pi(h_n(t) = a \mid g = 0) := \pi_{h_n(t)}(a \mid g = 0).$$

The Lipschitz continuity and positivity of  $u_n(t)$  and  $h_n(t)$  justify the use of Bayes' formula for probability density functions, and we get

$$\pi(h_n(t) = a \mid h_n(1) = 0) = \frac{\pi(h_n(t) = a) \pi(h_n(1) = 0 \mid h_n(t) = a)}{\pi(h_n(1) = 0)}. \tag{5.14}$$

Since  $h_n(1) - h_n(t)$  has the same distribution as  $h_n(1 - t)$ , we see by (5.14) that

$$\pi(h_n(t) = a \mid h_n(1) = 0) = c_n \pi(h_n(t) = a) \pi(h_n(1 - t) = -a). \tag{5.15}$$

We know that  $h_n(t)$  converges i.d. to a Gaussian random variable when  $n \rightarrow \infty$ . Namely, we see from (5.12) and the central limit theorem that  $\lim_{n \rightarrow \infty} h_n(t) = h(t)$ , where  $h(t)$  is Brownian motion with  $\mathbb{E} h(t) = 0$ ,  $\mathbb{E}(h(t) - h(s))^2 = |t - s| \sigma_p^2$  and  $t \geq s$ . This is not quite enough for our purposes and we need to modify the proof of the central limit theorem.

Denote the characteristic function of  $\xi_1$  by  $\varphi(s) = \mathbb{E} e^{is\xi_1}$ . Fourier transforming  $\exp(-|t|^p)$  shows that  $\varphi \in C^\infty(\mathbb{R})$ . Since  $\varphi''(0) = -\sigma_p^2$  by (5.6), there are such  $\sigma_0 < \sigma_p$  and  $\varepsilon > 0$  that

$$|\varphi(s)| \leq \exp(-\sigma_0^2 s^2 / 2) \quad \text{for } |s| < \varepsilon.$$

We write  $\varphi = \varphi_1 + \varphi_2$  with  $\varphi_1(s) = 0$  for  $|s| \geq \varepsilon$  and  $\varphi_2(s) = 0$  for  $|s| < \varepsilon$ . Further, we denote  $a = \sup |\varphi_2(s)| < 1$ .

It is well known that the characteristic function of the random variable  $S_j$  is  $\Psi_j(s) = (\varphi(s/\sqrt{j}))^j$ . By the central limit theorem,

$$\lim_{j \rightarrow \infty} \Psi_j(s) = \exp(-\sigma_p^2 |s|^2 / 2) \quad \text{for } s \in \mathbb{R}. \tag{5.16}$$

As the supports of  $\varphi_1$  and  $\varphi_2$  are disjoint, we see that

$$\Psi_j(s) = \Psi_j^1(s) + \Psi_j^2(s) = \left( \varphi_1 \left( \frac{s}{\sqrt{j}} \right) \right)^j + \left( \varphi_2 \left( \frac{s}{\sqrt{j}} \right) \right)^j. \tag{5.17}$$

Now we see that for any  $q \geq 1$

$$\|\Psi_j^2\|_{L^q}^q \leq a^{q(j-1)} \int_{\mathbb{R}} \left| \varphi_2 \left( \frac{s}{\sqrt{j}} \right) \right|^q ds \leq ca^{q(j-1)} j^{1/2} \rightarrow 0. \tag{5.18}$$

Moreover, we have the following estimate for  $\Psi_j^1$ :

$$|\Psi_j^1(s)| \leq \left( \exp \left( -\sigma_0^2 \left( \frac{s}{\sqrt{j}} \right)^2 / 2 \right) \right)^j = \exp(-\sigma_0 s^2 / 2). \tag{5.19}$$

Since  $\exp(-\sigma_0 s^2 / 2) \in L^q(\mathbb{R})$  for any  $q \geq 1$ , we see by (5.16)–(5.19) that

$$\lim_{j \rightarrow \infty} \|\Psi_j - \exp(-\sigma_p^2 | \cdot |^2 / 2)\|_{L^q(\mathbb{R})} = 0, \quad 1 \leq q < \infty. \tag{5.20}$$

By (5.13), the characteristic function of  $h_n(t)$  is

$$V_n(s) = \Psi_{\theta(n,t)}(k_{n,t}s) \cdot \varphi(r_{n,t}s). \tag{5.21}$$

Note that

$$\lim_{n \rightarrow \infty} \|\varphi(r_{n,t}s) - 1\|_{C^1(-L,L)} = 0 \quad \text{for any } L, |\varphi(r_{n,t}s)| \leq 1. \tag{5.22}$$

The characteristic function of  $h_n(1 - t)$ , denoted by  $G_n(s)$ , has a similar expression. Then by (5.15) the characteristic function of  $u_n(t)$  has the form

$$\Phi_n(s) = c_n (V_n * G_n)(s).$$

By (5.20)–(5.22) we see that  $(V_n * G_n)(s)$  converges for any  $s \in \mathbb{R}$ :

$$\lim_{n \rightarrow \infty} (V_n * G_n)(s) = (V * G)(s),$$

where

$$V(s) = \lim_{n \rightarrow \infty} V_n(s) \quad \text{and} \quad G(s) = \lim_{n \rightarrow \infty} G_n(s).$$

Since  $V(s)$  and  $G(s)$  are Gaussian functions, the condition  $\Phi_n(0) = 1$  implies that the normalization constants  $c_n = \pi(h_n(1) = 0)^{-1}$  converge to a positive constant when  $n \rightarrow \infty$ . Thus, using (5.20) and (5.21) we see that

$$\begin{aligned} \lim_{n \rightarrow \infty} \Phi_n(s) &= \Phi(s; t) := \exp(-\sigma(t)s^2/2), \\ \frac{1}{\sigma(t)^2} &= \frac{1}{t\sigma_p^2} + \frac{1}{(1-t)\sigma_p^2}. \end{aligned} \tag{5.23}$$

Since limit (5.23) exists at every  $s$  and the limit function  $\Phi(s; t)$  is continuous at  $s = 0$ , it follows from Levy’s continuity theorem [11, theorem 9.8.2] that there is such a random variable  $U_t$  that

$$\lim_{n \rightarrow \infty} \mathbb{E}(h_n(t) \mid h_n(1) = 0) = U_t \quad \text{i.d.}$$

and that the characteristic function of  $U_t$  is  $\Phi(s; t)$ . Claim (i) is proved for  $Q = 1$ .

To prove claim (ii) of the theorem we consider  $L^2$ -bounds. We see that

$$\mathbb{E}(|u_n|^2) = \int_{\mathbb{R}} s^2 \pi(u_n = s) ds = -\partial_s^2 \Phi_n(s)|_{s=0} = -c_n (\partial_s V_n) * (\partial_s G_n)(s)|_{s=0}.$$

In view of the definitions of  $V_n$  and  $G_n$ , let us consider  $\partial_s \Psi_j$ :

$$\partial_s \Psi_j(s) = \partial_s \left( \varphi \left( \frac{s}{\sqrt{j}} \right) \right)^j = j \frac{1}{\sqrt{j}} (\partial_s \varphi) \left( \frac{s}{\sqrt{j}} \right) \left( \varphi \left( \frac{s}{\sqrt{j}} \right) \right)^{j-1}. \tag{5.24}$$

Fourier transforming  $\exp(-|t|^p)$  shows that  $\varphi \in C^\infty(\mathbb{R})$  and that

$$\begin{aligned} |\varphi(s)| &\leq c(1 + |s|)^{-1-p}, & s \in \mathbb{R}, \\ \varphi(s) &= 1 - \frac{\sigma_p}{2} s^2 + O(s^3), & s \text{ near } 0. \end{aligned}$$

Thus  $|s^{-1} \partial_s \varphi(s)| \leq c'$  for  $s \in \mathbb{R}$  and we get the estimate

$$|\partial_s \Psi_j(s)| = \left| \frac{(\partial_s \varphi)(s/\sqrt{j})}{s/\sqrt{j}} \right| \left| s \left( \varphi \left( \frac{s}{\sqrt{j}} \right) \right)^{j-1} \right| \leq c' \left| s \left( \varphi \left( \frac{s}{\sqrt{j}} \right) \right)^{j-1} \right|.$$

Writing  $\varphi = \varphi_1 + \varphi_2$  as before we see that for  $1 \leq q < \infty$

$$\left\| s \left( \varphi_2 \left( \frac{s}{\sqrt{j}} \right) \right)^{j-1} \right\|_{L^q}^q \leq a^{q(j-3)} \int_{\mathbb{R}} s^q \left| \varphi_2 \left( \frac{s}{\sqrt{j}} \right) \right|^{2q} ds \leq ca^{q(j-3)} j^{(1+q)/2} \rightarrow 0$$

as  $j \rightarrow \infty$ . Moreover, we see that

$$\left| s \left( \varphi_1 \left( \frac{s}{\sqrt{j}} \right) \right)^{j-1} \right| \leq s \left( \exp \left( -\sigma_0^2 \left( \frac{s}{\sqrt{j}} \right)^2 / 2 \right) \right)^{j-1} \leq s \exp(-\sigma_0^2 s^2 / 4),$$

which is an integrable bound. Thus using (5.21) and (5.24) we see that

$$\begin{aligned} |V_n(s)| + |\partial_s V_n(s)| &\leq C_2 s \exp(-\sigma_0^2 s^2 / 8), & |s| < n^{1/2} \varepsilon, \\ \lim_{n \rightarrow \infty} \int_{\mathbb{R} \setminus [-n^{1/2} \varepsilon, n^{1/2} \varepsilon]} (|V_n(s)|^q + |\partial_s V_n(s)|^q) ds &= 0, & 1 \leq |q| < \infty. \end{aligned} \tag{5.25}$$

Thus we have on interval  $[-n^{1/2} \varepsilon, n^{1/2} \varepsilon]$  an uniform integrable bound and in the complement of this interval we can estimate  $L^q$ -norms uniformly.

Using (5.25) we see for  $V_n(s)$  and  $G_n(s)$  that

$$|(\partial_s V_n) * (\partial_s G_n)(0)| \leq \|\partial_s V_n\|_{L^2} \|\partial_s G_n\|_{L^2} \leq C_3.$$

Thus  $\mathbb{E}(|u_n(t)|^2) \leq C_4$ , and in particular, the family  $u_n(t)$  is uniformly integrable.

It remains to prove claim (i) for  $Q > 1$  and claim (iii). We prove them together by considering the joint distribution of  $u_n(t_k)$  and  $\langle A_j, u_n \rangle$  for  $j = 1, \dots, N$ . For this, we denote  $A_{N+k} = \delta(t - t_k)$  for  $k = 1, \dots, Q$  implying that

$$u_n(t_k) = \langle A_{N+k}, u_n \rangle, \quad j = 1, \dots, Q.$$

Denote

$$c_{n\ell}^j = \int_0^1 \psi_\ell^n(t) A_j(dt), \quad b_{nk}^j = \sum_{\ell=k}^{n+1} c_{n\ell}^j, \tag{5.26}$$

and write using (5.9), (5.11) and (5.26) and changing the order of summation

$$\langle A_j, h_n \rangle = \sum_{\ell=1}^{n+1} h_\ell^n c_{n\ell}^j = \sum_{\ell=1}^{n+1} \left( \frac{\tilde{\alpha}^{-1/p}}{(n+1)^{1/2}} \sum_{k=1}^{\ell} \xi_k^n \right) c_{n\ell}^j = \frac{\tilde{\alpha}^{-1/p}}{(n+1)^{1/2}} \sum_{k=1}^{n+1} b_{nk}^j \xi_k^n, \tag{5.27}$$

for  $j = 1, \dots, N + Q$ . Now  $Y_{nk}(\omega) := \tilde{\alpha}^{-1/p} (n+1)^{-1/2} (b_{nk}^j \xi_k^n(\omega))_{j=1}^{N+Q}$  are independent random vectors in  $\mathbb{R}^{N+Q}$  and

$$(\langle A_j, h_n \rangle)_{j=1}^{N+Q} = \sum_{k=1}^{n+1} Y_{nk}. \tag{5.28}$$

We prove that (5.28) converges in distribution to a Gaussian variable when  $n \rightarrow \infty$ . By the Fabian–Hannan version of the Lindenberg central limit theorem [13] it is enough to show that

$$\lim_{n \rightarrow \infty} \sum_{k=1}^{n+1} \mathbb{E}(|\beta Y_{nk}|^2 1_{\{|\beta Y_{nk}| > \varepsilon\}}) = 0 \tag{5.29}$$

for arbitrary  $\beta \in \mathbb{R}^{N+Q} \setminus 0$  and  $\varepsilon > 0$ . Here  $1_{\{|\beta Y_{nk}| > \varepsilon\}}$  is the indicator function equal to 1 if  $|\beta Y_{nk}| > \varepsilon$  and zero otherwise.

Let  $\beta \in \mathbb{R}^{N+Q} \setminus 0$  and  $\varepsilon > 0$ . Set  $\gamma_{nk} = \sum_{j=1}^{N+Q} \beta_j b_{nk}^j$  for  $k = 1, \dots, n + 1$ . Note that  $|b_{nk}^j| \leq \|A_j\|_{TV}$  and  $|\gamma_{nk}| \leq C|\beta|$ . Estimate

$$\begin{aligned} \sum_{k=1}^{n+1} \mathbb{E}(|\beta Y_{nk}|^2 1_{\{|\beta Y_{nk}| > \varepsilon\}}) &\leq \sum_{k=1}^{n+1} \mathbb{E} \left( \left| \frac{\tilde{\alpha}^{-1/p} \gamma_{nk} \xi_k^n}{(n+1)^{1/2}} \right|^2 1_{\{|\gamma_{nk} \xi_k^n| > \varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2}\}} \right) \\ &\leq \frac{c \tilde{\alpha}^{-2/p}}{n+1} \sum_{k=1}^{n+1} \gamma_{nk}^2 \int_{|\gamma_{nk} x| > \varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2}} x^2 e^{-|x|^p/2} dx \\ &\leq \frac{c \tilde{\alpha}^{-2/p}}{n+1} \sum_{k=1}^{n+1} \gamma_{nk}^2 \exp(-|\gamma_{nk}|^{-1} \varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2} / 3). \end{aligned}$$

Since  $|\gamma_{nk}|^{-1} \geq C^{-1} |\beta|^{-1}$  we see that sum in the limit (5.29) is defined and formula (5.29) is true. Hence  $(\langle A_j, h_n \rangle)_{j=1}^{N+Q}$  converges in distribution to a Gaussian variable. The limit is  $(\langle A_j, h \rangle)_{j=1}^{N+Q}$ , where  $h(t)$  is Brownian motion on  $t \in [0, 1]$ ,  $h(0) = 0$ .

We have proved pointwise convergence; next we consider convergence in  $L^q$ . To simplify notation we replace  $s''$  by the approximation

$$s_n = \frac{r_n}{n}, \quad r_n = \theta(n, s'') + 1.$$



The characteristic function of the random variable  $(\xi_k^n, b_{nk}^1 \xi_k^n, \dots, b_{nk}^{N+Q} \xi_k^n)$  is

$$\mathbb{E} \left( \exp \left( i \left( \xi_k^n \zeta + \sum_{j=1}^{N+Q} b_{nk}^j \xi_k^n \eta^j \right) \right) \right) = \varphi \left( \zeta + \sum_{j=1}^{N+Q} b_{nk}^j \eta^j \right).$$

Thus the characteristic function of  $(h(s_n), \langle A_1, h_n \rangle, \dots, \langle A_{N+Q}, h_n \rangle)$  is

$$\Phi_n(\zeta, \eta^1, \dots, \eta^{N+Q}) = \mathbb{E} \left( \exp \left( i \left( h_n(s_n) \zeta + \sum_{j=1}^{N+Q} a_n^j \eta^j \right) \right) \right) = \prod_{k=1}^{r_n} \varphi \left( \frac{\zeta + \sum_{j=1}^{N+Q} b_{nk}^j \eta^j}{\tilde{\alpha}^{1/p} (n+1)^{1/2}} \right),$$

where we used (5.27) and  $b_{nk}^j = 0$  for  $k > r_n$ . We have shown that  $(\langle A_j, h \rangle)_{j=1}^{N+Q}$  converges i.d. to a Gaussian variable. Thus there is such a Gaussian function  $\Phi(\zeta, \eta)$  that

$$\lim_{n \rightarrow \infty} \Phi_n(\zeta, \eta) = \Phi(\zeta, \eta) \tag{5.30}$$

for any  $\zeta \in \mathbb{R}$  and  $\eta = (\eta^1, \dots, \eta^{N+Q}) \in \mathbb{R}^{N+Q}$ .

Now, let  $\varepsilon, \sigma_2$  and  $a < 1$  be such that

$$\begin{aligned} |\varphi(t+s)| &\leq \exp(-\sigma_2|t+s|^2/2) && \text{for } |s| < \varepsilon, |t| \leq 2\varepsilon, \\ |\varphi(t+s)| &\leq a && \text{for } |s| < \varepsilon, |t| > 2\varepsilon. \end{aligned}$$

Fix  $\eta = (\eta^1, \dots, \eta^{N+Q})$  and set  $\beta = \sup_{n,k} |\sum_{j=1}^{N+Q} b_{nk}^j \eta^j|$ . Take  $n$  so large that  $|\beta| \tilde{\alpha}^{-1/p} (n+1)^{-1/2} < \varepsilon$ . Then

$$\begin{aligned} &1_{\{|\zeta| < 2\varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2}\}} \left| \prod_{k=1}^{r_n} \varphi \left( \frac{\zeta + \sum_{j=1}^{N+Q} b_{nk}^j \eta^j}{\tilde{\alpha}^{1/p} (n+1)^{1/2}} \right) \right| \\ &\leq 1_{\{|\zeta| < 2\varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2}\}} \exp \left( -\frac{\sigma_2}{2} \sum_{k=1}^{r_n} \left| \frac{\zeta + \sum_{j=1}^{N+Q} b_{nk}^j \eta^j}{\tilde{\alpha}^{1/p} (n+1)^{1/2}} \right|^2 \right) \\ &\leq \exp \left( \frac{\sigma_2}{\tilde{\alpha}^{1/p}} \left( \beta^2 - \frac{1}{6} |\zeta|^2 \right) \right) \end{aligned}$$

and

$$\begin{aligned} &\left\| 1_{\{|\zeta| > 2\varepsilon \tilde{\alpha}^{1/p} (n+1)^{1/2}\}} \prod_{k=1}^{r_n} \varphi \left( \frac{\zeta + \sum_{j=1}^{N+Q} b_{nk}^j \eta^j}{\tilde{\alpha}^{1/p} (n+1)^{1/2}} \right) \right\|_{L^q}^q \leq a^{q(r_n-1)} \int_{\mathbb{R}} \left| \varphi \left( \frac{t + \sum_{j=1}^N b_{n1}^j \eta^j}{\tilde{\alpha}^{1/p} (n+1)^{1/2}} \right) \right|^q dt \\ &\leq ca^{q(r_n-1)} \tilde{\alpha}^{1/p} (n+1)^{1/2} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This and (5.30) show that for any fixed  $\eta$

$$\lim_{n \rightarrow \infty} \|\Phi_n(\cdot, \eta) - \Phi(\cdot, \eta)\|_{L^q(\mathbb{R})} = 0, \quad 1 \leq q < \infty. \tag{5.31}$$

Consider next  $\vec{a}_n = (\langle A_j, u_n \rangle)_{j=1}^{N+Q}$  and  $\vec{b} = (b_1, \dots, b_{N+Q}) \in \mathbb{R}^{N+Q}$ . Then

$$\begin{aligned} R_n(f, b_1, \dots, b_{N+Q}) &= \pi(h_n(s_n) = f, \vec{a}_n = \vec{b} \mid h_n(1) = 0) \\ &= \frac{1}{\pi(h_n(1) = 0)} \pi(h_n(s_n) = f, \vec{a}_n = \vec{b}, h_n(1) = 0) \\ &= c_n \pi(h_n(1) = 0 \mid h_n(s_n) = f, \vec{a}_n = \vec{b}) \pi(h_n(s_n) = f, \vec{a}_n = \vec{b},) \\ &= c_n \pi(h_n(1) = 0 \mid h_n(s_n) = f) \pi(h_n(s_n) = f, \vec{a}_n = \vec{b},). \end{aligned}$$

Here in the last equality we have used the fact that  $h_j^n$  is a Markov sequence for  $j = 1, \dots, n$  and  $A_j$  are supported on  $[s', s'']$  with  $s'' < s_n$ , and thus

$$\pi(h_n(1) = 0 \mid h_n(s_n) = f, \vec{a}_n = \vec{b}) = \pi(h_n(1) = 0 \mid h_n(s_n) = f).$$

Let us now introduce an auxiliary variable  $d$  and a function

$$R_n^1(d, f, b_1, \dots, b_{N+Q}) = c_n \pi(h_n(1) = 0 \mid h_n(s_n) = f - d) \pi(h_n(s_n) = f, \vec{a}_n = \vec{b})$$

for which  $R_n(f, b_1, \dots, b_{N+Q}) = R_n^1(0, f, b_1, \dots, b_{N+Q})$ . Define functions  $W^n : \mathbb{R}^{1+N+Q} \rightarrow \mathbb{R}$  and  $K^n : \mathbb{R} \rightarrow \mathbb{R}$  by

$$W^n(f, \vec{b}) = \pi(h_n(s_n) = f, \vec{a}_n = \vec{b}),$$

$$K^n(f) = \pi(h_n(1) = 0 \mid h_n(s_n) = f) = \pi(h_n(1 - s_n) = -f),$$

and denote their Fourier transforms by  $\hat{W}^n(\xi, \eta)$  and  $\hat{K}^n(\xi)$ . If  $\zeta$  is the Fourier variable corresponding to  $d$ , the Fourier transform of  $R_n^1(d, f, b_1, \dots, b_{N+Q})$  is by (5.32)

$$\hat{R}_n^1(\zeta, \xi, \eta^1, \dots, \eta^{N+Q}) = c_n \hat{W}^n(\zeta, \eta^1, \dots, \eta^{N+Q}) \hat{K}^n(\xi + \zeta).$$

Then the characteristic function of the variable  $(u_n(s_n), \vec{a}_n)$  is

$$\Phi_n(\xi, \eta^1, \dots, \eta^{N+Q}) = c_n \int_{\mathbb{R}} R_n^1(\zeta, \xi, \eta^1, \dots, \eta^{N+Q}) d\zeta.$$

Since  $|\hat{W}^n(\zeta, \eta^1, \dots, \eta^{N+Q})| \leq 1$  and  $\hat{K}^n$  converge to a Gaussian function in  $L^1(\mathbb{R})$  by (5.16) and (5.31) we see that

$$\lim_{n \rightarrow \infty} \Phi_n(\xi, \eta^1, \dots, \eta^{N+Q}) = \left( \lim_{n \rightarrow \infty} c_n \right) \int_{\mathbb{R}} \left( \lim_{n \rightarrow \infty} R_n^1(\zeta, \xi, \eta^1, \dots, \eta^{N+Q}) \right) d\zeta.$$

Here,  $R_n^1$  converges to a Gaussian function. Thus the characteristic function  $\Phi_n(\xi, \eta)$  with fixed  $\xi$  and  $\eta$  converges to the Gaussian function  $\Phi(\xi, \eta)$ , which implies convergence of joint distributions i.d. □

### 5.2. Convergence of posterior distributions

Here we prove theorems 5.1 and 5.2.

Let  $v_n$  and  $v$  be random variables taking values in  $C(0, 1)$  and assume  $v_n \rightarrow v$  weakly i.d. By applying Skorohod's representation theorem (see e.g. [1]), and enlarging the probability space  $\Omega$  if necessary, we have

**Proposition 5.1.** *Let  $t \in (0, 1)$  be given. The random variables  $v_n(t, \omega), v(t, \omega), a_j^n = \langle v_n, A_j \rangle$ , and  $a_j = \langle v, A_j \rangle, j = 1, \dots, N$  have such versions  $\tilde{v}_n(t, \omega), \tilde{v}(t, \omega), \tilde{a}_j$  and  $\tilde{a}_j^n$  that almost surely*

$$\lim_{n \rightarrow \infty} \tilde{v}_n(t) = \tilde{v}(t), \quad \lim_{n \rightarrow \infty} \tilde{a}_j^n = \tilde{a}_j, \quad j = 1, \dots, N.$$

Using the versions  $\tilde{a}_j^n$  and  $\tilde{a}_j$  of  $\langle A_j, v_n \rangle$  and  $\langle A_j, v \rangle$ , respectively, we define random variables  $\tilde{m}_j^n = \tilde{a}_j^n + \varepsilon_j$  and  $\tilde{m}_j = \tilde{a}_j + \varepsilon_j$ . Here the errors  $\varepsilon_j \sim N(0, 1)$  are the same independent random variables.

**Proof** (theorem 5.1). Let  $F$  be either the identity map  $F(s) = s$  or  $F(s) = \chi_{(-\infty, \lambda]}(s)$  with  $\lambda \in \mathbb{R}$ . Since the random variables  $F(\tilde{v}_n(t))$  are uniformly integrable, converge a.s. (and thus i.p.) and the limiting variable  $F(\tilde{v}(t))$  is in  $L^1(\Omega)$ , we have by Vitali's convergence theorem [11, theorem 10.3.6] that

$$\lim_{n \rightarrow \infty} \|F(\tilde{v}_n(t)) - F(\tilde{v}(t))\|_{L^1(\Omega)} = 0.$$

Consider the random variables  $z_n = (\tilde{v}_n(t), \tilde{a}_1^n, \dots, \tilde{a}_N^n) = (z_n^0, z_n')$   $\in \mathbb{R}^{N+1}$  and  $z = (\tilde{v}(t), \tilde{a}_1, \dots, \tilde{a}_N) = (z^0, z') \in \mathbb{R}^{N+1}$ . Assume given a realization  $\hat{m} \in \mathbb{R}^N$  of the measurement and denote  $g(y, \hat{m}) = \pi_\varepsilon(y' - \hat{m}), y = (y^0, y') \in \mathbb{R}^{N+1}$ .

For clarity, we start our computations with the case where the laws  $P_{z_n}$  and  $P_z$  of  $z_n$  and  $z$  are absolutely continuous with respect to the Lebesgue measure in  $\mathbb{R}^{N+1}$  and there exist continuous probability density functions  $\pi_{z_n}(y)$  and  $\pi_z(y)$ .

Note that  $\tilde{m}^n$  has a smooth positive probability density function in  $\mathbb{R}^N$  given by

$$\pi(\tilde{m}^n = \hat{m}) = \int_{\mathbb{R}^{N+1}} \pi_\varepsilon(y' - \hat{m})\pi_{z_n}(y^0, y') \, dy = \int_{\mathbb{R}^{N+1}} g(y', \hat{m})\pi_{z_n}(y^0, y') \, dy. \tag{5.32}$$

A similar formula holds for  $\pi(\tilde{m} = \hat{m})$  and thus we have

$$\pi(\tilde{m}^n = \hat{m}) = \mathbb{E}(g(z'_n, \hat{m})), \quad \pi(\tilde{m} = \hat{m}) = \mathbb{E}(g(z', \hat{m})).$$

Since  $g$  is a bounded continuous function and  $z'_n \rightarrow z_n$  weakly, we see that

$$\lim_{n \rightarrow \infty} \pi(\tilde{m}^n = \hat{m}) = \pi(\tilde{m} = \hat{m}). \tag{5.33}$$

Moreover,  $\pi_{\tilde{m}^n}$  is a smooth function and we have by Bayes' formula

$$\begin{aligned} \pi(\tilde{v}_n(t) = z^0 | \tilde{m}^n = \hat{m}) &= \frac{1}{\pi(\tilde{m}^n = \hat{m})} \int_{\mathbb{R}^N} \pi_{z_n, \tilde{m}}(z^0, y', \hat{m}) \, dy', \\ &= \frac{1}{\pi(\tilde{m}^n = \hat{m})} \int_{\mathbb{R}^N} g(y', \hat{m})\pi_{z_n}(z^0, y') \, dy'. \end{aligned}$$

By (5.2),  $\tilde{E}(F(\tilde{v}_n(t)) | \tilde{m}^n = \hat{m})$  is equal to

$$\lim_{r \rightarrow 0} \frac{\int_{B(\hat{m}, r)} \int_{\mathbb{R}} F(y^0)\pi(\tilde{v}_n(t) = y^0, \tilde{m}^n = w) \, dy^0 \, dw}{\int_{B(\hat{m}, r)} \pi(\tilde{m}^n = w) \, dw} \tag{5.34}$$

and taking the integral over  $\mathbb{R}$  outside the limit and letting  $r \rightarrow 0$  leads to

$$\begin{aligned} \tilde{E}(F(\tilde{v}_n(t)) | \tilde{m}^n = \hat{m}) &= \int_{\mathbb{R}} F(y^0) \frac{\pi(\tilde{v}_n(t) = y^0, \tilde{m}^n = \hat{m})}{\pi(\tilde{m}^n = \hat{m})} \, dy^0 \\ &= \frac{1}{\pi(\tilde{m}^n = \hat{m})} \int_{\mathbb{R}^{N+1}} F(y^0)g(y, \hat{m})\pi_{z_n}(y^0, y') \, dy. \end{aligned} \tag{5.35}$$

Since a similar formula holds for  $u_B(t)$ , we have proved

$$\begin{aligned} \tilde{E}(F(\tilde{v}_n(t)) | \tilde{m}_n = \hat{m}) &= \frac{1}{\pi(\tilde{m}_n = \hat{m})} \mathbb{E}(F(z_n^0)g(z'_n, \hat{m})), \\ \tilde{E}(F(\tilde{v}(t)) | \tilde{m} = \hat{m}) &= \frac{1}{\pi(\tilde{m} = \hat{m})} \mathbb{E}(F(z^0)g(z', \hat{m})). \end{aligned} \tag{5.36}$$

In the general case, where the laws  $P_{z_n}$  and  $P_z$  of  $z_n$  and  $z$  are not absolutely continuous, we have to replace formula (5.32) by

$$\pi(\tilde{m}^n = \hat{m}) = \int_{\mathbb{R}^{N+1}} g(y, \hat{m})P_{z_n}(dy). \tag{5.37}$$

Again, since  $g$  is smooth by (5.37), we see that  $\pi_{\tilde{m}^n}$  is smooth. Also, we can replace formula (5.34) by

$$\tilde{E}(F(\tilde{v}_n(t)) | \tilde{m}^n = \hat{m}) = \lim_{r \rightarrow 0} \frac{\int_{B(\hat{m}, r)} \left( \int_{\mathbb{R}^N \times \mathbb{R}} F(y^0)\pi_\varepsilon(y' - w)P_{z_n}(dy) \right) \, dw}{\int_{B(\hat{m}, r)} \pi(\tilde{m}^n = w) \, dw}.$$

Since  $\pi(\tilde{m}^n = w)$  is smooth, there is  $C > 0$  such that

$$\left| \frac{\int_{B(\hat{m}, r)} \pi_\varepsilon(y' - w) \, dw}{\int_{B(\hat{m}, r)} \pi(\tilde{m}^n = w) \, dw} \right| \leq C \quad \text{for } y' \in \mathbb{R}^N.$$

Thus Fubini’s theorem and Lebesgue’s theorem of dominated convergence give an analogue of formula (5.35):

$$\tilde{E}(F(\tilde{v}_n(t))|\tilde{m}^n = \hat{m}) = \frac{1}{\pi(\tilde{m}^n = \hat{m})} \int_{\mathbb{R}^{N+1}} F(y^0)g(y', \hat{m})P_{z_n}(dy).$$

These formulae imply (5.33) and (5.36) also in the general case.

Let  $H(y) = F(y^0)g(y', \hat{m})$ . Since  $F(z_n^0) \rightarrow F(z^0)$  in  $L^1(\Omega)$  and  $|g| \leq 1$ , we have  $\lim_{n \rightarrow \infty} H(z_n) = H(z)$  in  $L^1(\Omega)$ . This and (5.33) imply that

$$\lim_{n \rightarrow \infty} \tilde{E}(F(\tilde{v}_n(t))|\tilde{m}^n = \hat{m}) = \tilde{E}(F(\tilde{v}(t))|\tilde{m} = \hat{m}).$$

Since the distributions of  $v_n(t)$  and  $\tilde{v}_n(t)$  as well as those of  $m^n$  and  $\tilde{m}^n$  coincide, we have by (5.2)

$$\tilde{E}(F(v_n(t))|m^n = \hat{m}) = \tilde{E}(F(\tilde{v}_n(t))|\tilde{m}^n = \hat{m}).$$

This proves the assertion. □

**Proof** (theorem 5.2). Let us first consider the case  $\alpha_n = \tilde{\alpha}(n + 1)^{1-p/2}$ . By theorem 5.3,  $u_n \rightarrow u_B$  weakly i.d.,  $u_n(t), t \in (0, 1)$ , are uniformly integrable and  $u_B(t) \in L^1(\Omega)$ . Then (5.3) and (5.4) follow from theorem 5.1.

When  $\alpha_n = \tilde{\alpha}(n + 1)^q$ , we see that  $w_n(t) = \tilde{\alpha}(n + 1)^{(q-1-p/2)/p}\tilde{u}_n(t)$  converges to the Brownian bridge. Thus  $\tilde{u}_n(t)$  converges to zero in  $L^1(\Omega)$  when  $q < 1 - p/2$ . When  $q > 1 - p/2$ , we see that  $u_n(t)$  cannot converge even i.d. □

## 6. Computational results

### 6.1. The model problem


Charge coupled devices (CCD) are commonly used in digital cameras and medical x-ray imaging devices. CCDs typically consist of a two-dimensional array of pixels capable of measuring the amount of visible light illuminating the area of the pixel over a period of time. We give a rough model for the measurement of the intensity distribution of light on one row of CCD pixels.

We take the quantity  $u$  in (1.1) to be a real-valued function on the unit interval  $[0, 1]$ . Given  $N > 1$ , we divide the subinterval  $[\frac{1}{N+2}, \frac{N+1}{N+2}] \subset [0, 1]$  into  $N$  pixels  $[x_j^{N+1}, x_{j+1}^{N+1}]$  with  $x_j^{N+1} = j/(N + 2)$  for  $j = 1, \dots, N$ . The measurement of the  $j$ th pixel is

$$m_j = \langle A_j, u \rangle + \varepsilon_j = \int_{x_j^{N+1}}^{x_{j+1}^{N+1}} u(t) dt + \varepsilon_j, \tag{6.1}$$

where  $A_j = \chi_{(x_j^{N+1}, x_{j+1}^{N+1})}$  as shown in figure 1, and  $\varepsilon_j$  are normally distributed random numbers with standard deviation  $\sigma > 0$ . The numbers  $\varepsilon_j$  model measurement errors resulting from quantum and electronic noise of the CCD.

We use the  $p$ -variation distribution in  $Y_n$  with  $p = 1$  or  $p = 2$  for representing *a priori* information on  $u$ . For convenience, we take the number of pixels to be of the form  $N = 2^L - 2$  with  $L > 1$ , and the dimension  $n = 2^\ell - 1$  is chosen to be greater than the number of measurements:  $\ell > L$ . In this case the grid  $x_1^{N+1}, \dots, x_N^{N+1}$  is a subset of the grid  $x_0^n, \dots, x_{n+1}^n$ .



**Figure 1.** Idealized one-dimensional model for the measurement. The function represents a distribution of light on the interval  $[0, 1]$ . Pixels are represented by intervals and measurements are integrals of light intensity over those intervals. No measurement is made on the leftmost and rightmost intervals.

6.2. Computational methods

6.2.1. Computation of MAP estimate with  $p = 2$ . Denote  $U = [u_1^n, \dots, u_n^n]^T$ . Consider the minimization problem

$$\tilde{U} = \arg \min_{U \in \mathbb{R}^n} \left( \|AU - \hat{m}\|_{\mathbb{R}^N}^2 + \frac{2\sigma^2\alpha_n}{\Delta x^n} \|\mathcal{D}U\|_{\mathbb{R}^{n+1}}^2 \right), \tag{6.2}$$

where  $\Delta x^n = (n + 1)^{-1}$ . The  $N \times n$  matrix  $\mathcal{A}$  implements the measurement:

$$(\mathcal{A}U)_k = \left\langle A_k, \sum_{j=1}^n u_j^n \psi_j^n \right\rangle, \quad k = 1, 2, \dots, N, \tag{6.3}$$

where the roof-top basis functions  $\psi_j^n$  for the space  $Y_n$  are as in definition 2.1. Integration of the piecewise linear functions in (6.3) over the intervals  $[x_j^{N+1}, x_{j+1}^{N+1}]$  is implemented simply and exactly by the classical trapezoidal rule. Thus the  $j$ th row of  $\mathcal{A}$  takes the form

$$\left[ \underbrace{0, \dots, 0}_{j2^{l-L}-1}, \frac{1}{2} \Delta x^n, \underbrace{\Delta x^n, \Delta x^n, \dots, \Delta x^n}_{2^{l-L}-1}, \frac{1}{2} \Delta x^n, 0, 0, \dots, 0 \right].$$

Prior information is coded into  $\mathcal{D}$ , the  $(n + 1) \times n$  matrix defined by

$$(\mathcal{D}U)_k = u_k^n - u_{k-1}^n, \quad k = 1, \dots, n + 1, \quad u_0^n = 0 = u_{n+1}^n.$$

Following Varah [34] we write (6.2) in stacked form

$$\begin{bmatrix} \mathcal{A} \\ \left(\frac{2\sigma^2\alpha_n}{\Delta x^n}\right)^{1/2} \mathcal{D} \end{bmatrix} U = \begin{bmatrix} \hat{m} \\ 0 \end{bmatrix} \tag{6.4}$$

and compute  $\tilde{U} = [\tilde{u}_1^n, \dots, \tilde{u}_n^n]^T$  as the least-squares solution of (6.4) using the Moore–Penrose pseudoinverse. In view of (2.2), (4.1), (4.5) and (6.2) we have

$$u_n^{\text{MAP}}(t; 2, \alpha_n) = \sum_{j=1}^n \tilde{u}_j^n \psi_j^n(t).$$

6.2.2. Computation of MAP estimate with TV prior. Consider the non-unique minimization problem

$$\tilde{U} \in \arg \min_{U \in \mathbb{R}^n} \left( \frac{1}{2\sigma^2} \|AU - \hat{m}\|_{\mathbb{R}^N}^2 + \alpha_n \sum_{j=1}^{n+1} |(\mathcal{D}U)_j| \right), \tag{6.5}$$

where the matrices  $\mathcal{A}$  and  $\mathcal{D}$  are as in section 6.2.1. We write  $\mathcal{D}U$  in the form

$$V_+, V_- \in \mathbb{R}_+^{n+1}, \quad V_+ - V_- = \mathcal{D}U, \tag{6.6}$$

where  $\mathbb{R}_+ := \{x \in \mathbb{R} : x \geq 0\}$ . Now problem (6.5) is equivalent to

$$\tilde{U} = \arg \min \left( \frac{1}{2\sigma^2} \|\mathcal{A}U\|_2^2 - \frac{1}{\sigma^2} \hat{m}^T \mathcal{A}U + \alpha_n \mathbf{1}^T V_+ + \alpha_n \mathbf{1}^T V_- \right), \tag{6.7}$$

where  $\tilde{U} = [\tilde{U}^T \tilde{V}_+^T \tilde{V}_-^T]^T$  and  $\mathbf{1} = [1, 1, \dots, 1]^T \in \mathbb{R}^{n+1}$ ; the minimum is taken over  $U \in \mathbb{R}^n, V_+, V_- \in \mathbb{R}^{n+1}$  satisfying (6.6). The solution of (6.7) with constraints (6.6) satisfies

$$(\tilde{V}_+)_j = \max((\mathcal{D}\tilde{U})_j, 0), \quad (\tilde{V}_-)_j = \max((-\mathcal{D}\tilde{U})_j, 0).$$

Write now  $\mathcal{U} = [U^T V_+^T V_-^T]^T$  and

$$H = \begin{bmatrix} \frac{1}{\sigma^2} \mathcal{A}^T \mathcal{A} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad f = \begin{bmatrix} -\frac{1}{\sigma^2} \mathcal{A}^T \hat{m} \\ \alpha_n \mathbf{1} \\ \alpha_n \mathbf{1} \end{bmatrix}.$$

Now problem (6.5) is converted to a standard quadratic minimization problem

$$\tilde{U} = \arg \min \left\{ \frac{1}{2} \mathcal{U}^T H \mathcal{U} + f^T \mathcal{U} \right\} \tag{6.8}$$

with linear constraints (6.6). We assume that we can find algorithmically an approximation to one of the possibly many solutions to (6.8) and (6.6). In view of (2.2), (4.1), (4.5) and (6.5) we define

$$u_n^{\text{MAP}}(t; 1, \alpha_n) = \sum_{j=1}^n \tilde{u}_j^n \psi_j^n(t).$$

6.2.3. *Computation of CM estimates.* Monte Carlo Markov chain (MCMC) methods can be used to generate a collection  $U^{(1)}, \dots, U^{(K)} \in \mathbb{R}^n$  of samples asymptotically distributed according to the posterior distribution

$$\tilde{\pi}(U | \hat{m}) = \tilde{c}_{n,p} \exp \left( -\frac{1}{2\sigma^2} \|\mathcal{A}U - \hat{m}\|_{\mathbb{R}^n}^2 - \frac{\alpha_n}{(n+1)^{1-p}} \sum_{v=1}^{n+1} |(\mathcal{D}U)_v|^p \right). \tag{6.9}$$

If  $K$  is large, we have


$$\int_{\mathbb{R}^n} U \tilde{\pi}(U | \hat{m}) dU \approx \frac{1}{K - k_0} \sum_{k=k_0+1}^K U^{(k)} =: \tilde{U}, \tag{6.10}$$

where the first  $k_0 > 0$  samples have been discarded because MCMC algorithms typically need such a *burn-in period* before the samples start to explore the posterior distribution representatively.


We use the Metropolis–Hastings (MH) algorithm [14, 16]. To implement the MH algorithm we define the *proposal distribution*  $Q(V, \cdot)$  on  $\mathbb{R}^n$ , parametrized by  $V \in \mathbb{R}^n$ , as follows. Fix  $1 \leq N_{\text{update}} \leq n$  and  $\kappa > 0$ . Pick randomly  $N_{\text{update}}$  distinct numbers from the set  $1, 2, \dots, n$  according to uniform probability distribution. Order the numbers and denote them by  $j_1, j_2, \dots, j_{N_{\text{update}}}$ . Then a candidate vector  $U \in \mathbb{R}^n$  is picked according to  $Q(V, \cdot)$  if  $U = V + \mathcal{E}_\kappa$ , where

$$\mathcal{E}_\kappa = [0, \dots, 0, \varepsilon'_{j_1}, 0, \dots, 0, \varepsilon'_{j_2}, 0, \dots, 0, \varepsilon'_{N_{\text{update}}}, 0, \dots, 0]^T$$

with  $\varepsilon'_{j_\ell} \sim N(0, \kappa)$  independent random numbers. Note that, if  $\pi_Q(V, U)$  denotes the density of  $Q(V, \cdot)$ , the *transition probabilities* are symmetric:  $\pi_Q(V, U) = \pi_Q(U, V)$ .



**Figure 2.** Left: simulated intensity distribution  $u(t)$ . Right: simulated noisy measurement  $\hat{m}$ . The dots are plotted at centre points of pixels.



**Figure 3.** Gaussian MAP estimates with three different choices of  $\alpha_{63}$ . The function  $u(t)$  is plotted with a thin line. Left: too small  $\alpha_{63}$ . Middle: satisfactory solution. Right: too large  $\alpha_{63}$ .

Due to the above symmetry, the MH algorithm takes the simple form

- (1) Set  $k := 0$  and initialize  $U^{(0)}$  by e.g.  $U^{(0)} := [0, \dots, 0]^T$ .
- (2) Set  $U := U^{(k)} + \mathcal{E}_\kappa$ .
- (3) If  $\tilde{\pi}(U | \hat{m}) \geq \tilde{\pi}(U^{(k)} | \hat{m})$  then set  $U^{(k+1)} = U$  and go to 5.
- (4) Draw a random number  $s$  from the uniform distribution on  $[0, 1]$ . If  $s \leq \frac{\tilde{\pi}(U | \hat{m})}{\tilde{\pi}(U^{(k)} | \hat{m})}$  then set  $U^{(k+1)} := U$ ; else set  $U^{(k+1)} := U^{(k)}$ .
- (5) If  $k = K$  then stop; else set  $k \leftarrow k + 1$  and go to 2.

We close this section by defining the *acceptance rate* of the Markov chain produced by the MH algorithm (discarding  $k_0$  first samples):

$$r = \frac{\text{number of accepted candidates}}{K - k_0}. \tag{6.11}$$

### 6.3. Results

In our numerical examples we take  $u$  to be the step function satisfying  $u(t) = 1$  for  $t \in [1/3, 2/3]$  and  $u(t) = 0$  otherwise. We consider a measurement with  $N = 2^5 - 2 = 30$  pixels and random errors  $\varepsilon_j$  with standard deviation  $\sigma = 0.001$ . See figure 2 for a plot of a realization of the measurement.


We perform all the computations with Matlab 6.5 running in a desktop PC computer equipped with a 2.8 GHz Intel Pentium 4 processor and 1 GB of RAM.

*6.3.1. The Gaussian case.* We start by determining a suitable parameter  $\alpha_{63}$  for MAP estimates with 2-variation prior and fixed discretization level  $\ell = 6, n = 63$ . See figure 3 for the least-squares solutions of (6.4) with various parameters. Based on visual inspection, we choose  $\alpha_{63} = 10$ .

We turn to computing MAP estimates with varying levels of discretization. We solve (6.4) with  $n = 63, 127, 255, 511, 1023, 2047, 4095$ , with  $\alpha_n = 10$  for all  $n$ . The solutions agree with good precision at the coarsest discretization level:

$$\max_{j=1, \dots, 63} \{|u_{63}^{\text{MAP}}(x_j^{63}; 2, \tilde{\alpha}_2) - u_n^{\text{MAP}}(x_j^{63}; 2, \tilde{\alpha}_2)|\} \leq 0.003. \tag{6.12}$$

The computation takes less than a second for  $n < 512$  and 295 s for  $n = 4095$ .



**Figure 4.** In all the plots in this figure, the coordinate axis limits are the same to allow easy comparison. Left column: MAP estimates for the TV prior with parameter  $\alpha_n = 135$  (thin line) and  $\alpha_n = 16.875\sqrt{n+1}$  (thick line). Right column: CM estimates for the TV prior with parameter  $\alpha_n = 135$  (thin line) and  $\alpha_n = 16.875\sqrt{n+1}$  (thick line).

**6.3.2. MAP estimates for the TV prior.** We determine a suitable parameter  $\alpha_{63}$  for MAP estimates with the TV prior at fixed discretization level  $\ell = 6, n = 63$  by numerical experimentation. As the result we choose  $\alpha_{63} = 135$ .

We compare two ways of choosing the parameter as a function of  $n$ , both satisfying  $\alpha_{63} = 135$ :

$$(i) \quad \alpha_n = 135, \quad (ii) \quad \alpha_n = 16.875\sqrt{n+1}. \quad (6.13)$$

We use MOSEK Optimization Toolbox's quadprog routine (available from [www.mosek.com](http://www.mosek.com)) to solve (6.8) with the constraints (6.6) in dimensions  $n = 255, 1023, 4095$ . Each computation takes less than 60 s of CPU time. See the left column in figure 4 for plots of the MAP estimates with choices (i) and (ii).

We note that small changes in parameter values changed the computation considerably, sometimes even resulting in divergence of the algorithm. We presume this is due to the non-uniqueness of the solution of the optimization problem. However, the presented results did not exhibit these problems and we believe them to be approximations to some functions in the set of solutions to the optimization problem.

**6.3.3. CM estimates for TV prior.** We compute CM estimates using the MH algorithm for  $n = 63, 255, 1023, 4095$ . Choices (i) and (ii) of  $\alpha_n$  given in (6.13) are compared. Parameters of the MCMC computations are given in table 1; in each case we take the zero vector as the initial guess. See the right column in figure 4 for plots of the CM estimates.



**Table 1.** Parameters of MCMC computations. The number  $n$  is the dimension of the problem,  $K - k_0$  is the number of samples used for computing the CM estimate,  $r$  is the acceptance rate defined in (6.11),  $N_{\text{update}}$  and  $\kappa$  are parameters of the proposal distribution and the last column indicates how many CPU hours the computations took.

$n$	$\alpha_n$	$K - k_0$	$r$	$N_{\text{update}}$	$\kappa$	Time (h)
63	135	80 000 000	0.23	10	0.082	3
255	135	80 000 000	0.24	10	0.082	6
255	270	80 000 000	0.24	10	0.041	6
1023	135	80 000 000	0.25	10	0.082	18
1023	540	80 000 000	0.25	10	0.021	16
4095	135	160 000 000	0.24	100	0.024	153
4095	1080	610 000 000	0.23	100	0.004	520

We actually use the MH algorithm slightly differently than explained in section 6.2.3. Denote by  $r_{1000}$  the *local acceptance rate* of the last 1000 samples. Choosing too large  $\kappa$  to start with leads to  $r_{1000} = 0$  and the chain does not move. On the other hand, choosing a very small  $\kappa$  results in a positive  $r_{1000}$  that, however, keeps growing until it reaches a value close to 1; then the candidates are always accepted and the chain moves very slowly. To overcome this problem we introduce automatic doubling of  $\kappa$  whenever  $r_{1000} > 0.35$ , but then the resulting chain is not Markov. However, after running for a while,  $r_{1000}$  becomes nearly constant and  $\kappa$  is not changed any more. An interpretation of our strategy is that we use the  $\kappa$ -doubling scheme to find a good initial guess for the  $K - k_0$  samples in the end of the chain that were drawn with constant  $\kappa$ . Those  $K - k_0$  most recent samples form a Markov chain.

#### 6.4. Discussion


We have computed the following statistical estimates for the posterior distribution of the model problem with varying levels of discretization:

$p$	Estimate	$\alpha_n$	Limit function
2	MAP	10	$\tilde{u}(\cdot; 2, 10)$
2	CM	10	$\tilde{u}(\cdot; 2, 10)$
1	MAP	135	$\tilde{u}(\cdot; 1, 135)$
1	MAP	$16.875\sqrt{n+1}$	0
1	CM	135	Does not exist
1	CM	$16.875\sqrt{n+1}$	Smooth

The column ‘limit function’ above indicates the expected result of each computation in light of theorems 4.1 and 5.2; the function  $\tilde{u}$  is defined by (4.2). How well do our computations agree with the theory?

The Gaussian computations in section 6.3.1, together with the error estimate (6.12), illustrate the convergence of the Gaussian MAP estimates (and CM estimates as well, since the two coincide in the Gaussian case).

In case of the TV MAP estimates, the choice  $\alpha_n = 16.875\sqrt{n+1}$  gives the zero estimate at the limit  $n \rightarrow \infty$ . This is evident from figure 4. Choosing  $\alpha_n = 135$  should lead to convergence to a limit function  $\tilde{u}(\cdot; 1, 135)$ . This is clear from the superposition of those estimates for  $n = 63, 255, 1023, 4095$  shown in figure 5.



**Figure 5.** Left: superposition of TV MAP estimates with  $n = 63, 255, 1023, 4095$  and choice (i), or  $\alpha_n = 135$ . Right: superposition of the TV CM estimates with  $n = 63, 255, 1023, 4095$  and choice (ii), or  $\alpha_n = 16.875\sqrt{n+1}$ .

The right column in figure 4 shows the CM estimates for the TV prior. We can see that the choice  $\alpha_n = 135$  leads to more and more oscillatory, divergent CM estimates, as expected (although the plot with  $n = 4095$  is not to be completely trusted due to the possibly insufficient number of samples used). On the other hand, the CM estimates with choice  $\alpha_n = 16.875\sqrt{n+1}$  are supposed to converge to a limit function. As the superposition in figure 5 reveals, the CM estimate for  $n = 4095$  is not of best possible quality. This is due to the very slow convergence of the chain; the computation took 520 h. However, in our view the degree of convergence is enough to conclude that the limit function is not edge-preserving.

### Acknowledgments

This work was supported by the Academy of Finland (no 72434 and 102175), the Finnish Technology Agency (TEKES), Instrumentarium Corporation, Imaging Division, Finland, and Invers Ltd, Finland. The second author was funded in part by the Grant-in-Aid for JSPS Fellows (no 0002757) of the Japan Society for the Promotion of Science. The authors are grateful for fruitful discussions with Jean Bricmont, Jari Kaipio, Ville Kolehmainen, Antti Kupiainen, Sari Lasanen, Geoff Nicholls, Teemu Pennanen, Erkki Somersalo and Johanna Tamminen. Finally, the authors thank Markku Lehtinen for pointing out the most essential observation that total variation priors converge to a Gaussian variable.

### References

- [1] Arcudi O 1998 Convergence of conditional expectations given the random variables of a Skorohod representation *Stat. Probab. Lett.* **40** 1–8
- [2] Attouch H and Wets R 1987 Epigraphical analysis. Analyse non lineaire (Perpignan, 1987) *Ann. Inst. Henri Poincaré Anal. Non Linéaire* **6** 73–100
- [3] Attouch H 1984 Variational convergence for functions and operators *Applicable Mathematics Series* (Boston, MA: Pitman) xiv+423 pp
- [4] Aubert G, Barlaud M, Blanc-Fraud M L and Charbonnier P 1997 Deterministic edge-preserving regularization in computed imaging *IEEE Trans. Image Process.* **5** 298–311
- [5] Aubert G and Kornprobst P 2000 *Mathematical Problems in Image Processing (Applied Mathematical Sciences vol 147)* (Berlin: Springer)
- [6] Boylan E 1971 Equiconvergence of martingales *Ann. Math. Stat.* **42** 552–9
- [7] Brascamp H J, Lieb E H and Lebowitz J L 1976 The statistical mechanics of anharmonic lattices *Proc. 40th Session of the International Statistical Institute (Warsaw, 1975)* vol 1 Invited papers *Bull. Inst. Int. Statist.* **46** 393–404
- [8] Demengel F and Temam R 1984 Convex functions of a measure and applications *Indiana Univ. Math. J.* **33** 673–709
- [9] Dobson D C and Santosa F 1996 Recovery of blocky images from noisy and blurred data *SIAM J. Appl. Math.* **56** 1181–98

- [10] Duoandikoetxea J 2001 *Fourier Analysis (Graduate Studies in Mathematics vol 29)* (Providence, RI: American Mathematical Society) xviii+222 pp
- [11] Dudley R M 1989 *Real Analysis and Probability (Mathematics Series)* (Pacific Grove, CA: Wadsworth & Brooks/Cole) xii+436 pp
- [12] Evans S N and Stark P B 2002 Inverse problems as statistics *Inverse Problems* **18** R55–97
- [13] Fabian V and Hannan J 1985 *Introduction to Probability and Mathematical Statistics* (New York: Wiley)
- [14] Gilks W R, Richardson S and Spiegelhalter D J 1996 *Markov Chain Monte Carlo in Practice* (London: Chapman and Hall)
- [15] Haario H, Laine M, Lehtinen M, Saksman S and Tamminen J 2004 MCMC methods for high dimensional inversion in remote sensing *J. R. Stat. Soc. Ser. B* **66** 591–607
- [16] Hastings W K 1970 Monte Carlo sampling methods using Markov chains and their applications *Biometrika* **57** 97–109
- [17] Kaipio J P, Kolehmainen V, Somersalo E and Vauhkonen M 2000 Statistical inversion and Monte Carlo sampling methods in electrical impedance tomography *Inverse Problems* **16** 1487–522
- [18] Kaipio J P, Kolehmainen V, Vauhkonen M and Somersalo E 1999 Inverse problems with structural prior information *Inverse Problems* **15** 713–29
- [19] Kolehmainen V, Siltanen S, Järvenpää S, Kaipio J P, Koistinen P, Lassas M, Pirttilä J and Somersalo E 2003 Statistical inversion for x-ray tomography with few radiographs: II. Application to dental radiology *Phys. Med. Biol.* **48** 1465–90
- [20] Lasanen S 2002 Discretizations of generalized random variables with applications to inverse problems *Dissertation University of Oulu. Ann. Acad. Sci. Fenn. Math. Diss.* No 130, 64 pp
- [21] Lehtinen M 2000 Personal communications
- [22] Lehtinen M S, Päivärinta L and Somersalo E 1989 Linear inverse problems for generalised random variables *Inverse Problems* **5** 599–612
- [23] D’Ambrogi B, Mäenpää S and Markkanen M 1999 Discretization independent retrieval of atmospheric ozone profile *Geophysica* **35** 87–99
- [24] Meyer Y 2001 *Oscillating Patterns in Image Processing and Nonlinear Evolution Equations (University Lecture Series vol 22)* (Providence, RI: American Mathematical Society)
- [25] Mosegaard K and Sambridge M 2002 Monte Carlo analysis of inverse problems *Inverse Problems* **18** R29–54
- [26] Naddaf A and Spencer T 1997 On homogenization and scaling limit of some gradient perturbations of a massless free field *Commun. Math. Phys.* **183** 55–84
- [27] Oksendal B 1998 *Stochastic Differential Equations. An Introduction with Applications (Universitext)* (Berlin: Springer) xx+324 pp
- [28] Rockafellar R T and Wets R J-B 1998 Variational analysis *Grundlehren der Mathematischen Wissenschaften vol 317* (Berlin: Springer) xiv+733 pp
- [29] Rudin L I, Osher S and Fatemi E 1992 Nonlinear total variation based noise removal algorithms *Physica D* **60** 259–68
- [30] Siltanen S, Kolehmainen V, Järvenpää S, Kaipio J P, Koistinen P, Lassas M, Pirttilä J and Somersalo E 2003 Statistical inversion for x-ray tomography with few radiographs: I. General theory *Phys. Med. Biol.* **48** 1437–63
- [31] Tamminen J 1999 *MCMC Methods for Inverse Problems (Geophysical Publications vol 48)* (Helsinki: Finnish Meteorological Institute)
- [32] Tikhonov A N 1963 Solution of incorrectly formulated problems and the regularization method *Sov. Math.–Dokl.* **4** 1035–8
- [33] Vakhania N, Tarieladze V I and Chobanyan S A 1987 *Probability Distributions on Banach Spaces (Mathematics and its Applications (Soviet Series) vol 14)* (Dordrecht: Reidel) xxvi+482 pp
- [34] Varah J M 1979 A practical examination of some numerical methods for linear discrete ill-posed problems *SIAM Rev.* **21** 100–11
- [35] Vogel C and Oman M 1998 Fast, robust total variation-based reconstruction of noisy, blurred images *IEEE Trans. Image Process.* **7** 813–24
- [36] Ziemer W 1989 *Weakly Differentiable Functions. Sobolev Spaces and Functions of Bounded Variation (Graduate Texts in Mathematics vol 120)* (Berlin: Springer) xvi+308 pp